

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/55, 9/14, C07K 16/40, C12Q 1/34, C12N 11/00	A1	(11) International Publication Number: WO 98/31816 (43) International Publication Date: 23 July 1998 (23.07.98)
(21) International Application Number: PCT/US98/00944 (22) International Filing Date: 16 January 1998 (16.01.98) (30) Priority Data: 60/035,404 17 January 1997 (17.01.97) US (71) Applicant (for all designated States except US): REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; 100 Church Street, S.E., Minneapolis, MN 55455 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): WACKETT, Lawrence, P. [US/US]; 1575 Merrill Street, St. Paul, MN 55108 (US). SADOWSKY, Michael, J. [US/US]; 710 Lovell Avenue, Roseville, MN 55113 (US). De SOUZA, Mervyn, L. [IN/US]; 1029 Raymond Avenue #10, St. Paul, MN 55114-11334 (US). MINSHULL, Jeremy, S. [GB/US]; 11 Homer Lane, Menlo Park, CA 94025 (US). (74) Agent: MCCORMACK, Myra, H.; Muetting, Raasch & Gebhardt, P.A., P.O. Box 581415, Minneapolis, MN 55458-1415 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: DNA MOLECULES AND PROTEIN DISPLAYING IMPROVED TRIAZINE COMPOUND DEGRADING ABILITY		
(57) Abstract This invention relates to the identification of homologs of atrazine chlorohydrolase and the use of these homologs to degrade s-triazine-containing compounds. In particular, this invention includes the identification of homologs of atrazine chlorohydrolase encoded by a DNA fragment having at least 95 % homology to the sequence from the nucleic acid-sequence beginning at position 236 and ending at position 1655 of SEQ ID NO:1, where the DNA fragment is capable of hybridizing under stringent conditions to SEQ ID NO:1 and has altered catalytic activity as compared with wild-type atrazine chlorohydrolase.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DNA MOLECULES AND PROTEIN DISPLAYING IMPROVED TRIAZINE COMPOUND DEGRADING ABILITY

Background of the Invention

5 More than 8 million organic compounds are known and many are
thought to be biodegradable by microorganisms, the principle agents for
recycling organic matter on Earth. In this context, microbial enzymes represent
the greatest diversity of novel catalysts. This is why microbial enzymes are
predominant in industrial enzyme technology and in bioremediation, whether
10 used as purified enzymes or in whole cell systems.

 There is increased interest in engineering bacterial enzymes for
improved industrial performance. For example, site directed mutagenesis of
subtilisin has resulted in the development of enzyme variants with improved
properties for use in detergents. Most applied enzymes, particularly those used
15 in biodegrading pollutants, however, are naturally evolved. That is, they are
unmodified from the form in which they were originally present in a soil
bacterium.

 For example, most bioremediation is directed against petroleum
hydrocarbons, pollutants that are natural products and thus have provided
20 selective pressure for bacterial enzyme evolution over millions of years.
Synthetic compounds not resembling natural products are more likely to resist
biodegradation and hence accumulate in the environment. This changes over a
bacterial evolutionary time scale; compounds considered to be
non-biodegradable several decades ago, for example PCBs and
25 tetrachloroethylene, are now known to biodegrade. This is attributed to recent
evolution and dispersal of the newly evolved gene(s) throughout microbial
populations by mechanisms such as conjugative plasmids and transposable DNA
elements.

 A better understanding of the evolution of new biodegradative
30 enzymes will reveal how nature cleanses the biosphere. Furthermore, the ability
to emulate the process in the laboratory may shave years off the lag period
between the introduction of a new molecular compound into the environment
and the development of a dispersed microbial antidote that will remove it.

Atrazine [2-chloro-4-(ethylamino)-6-(isopropylamino)-1,3,5-triazine]] is a widely used *s*-triazine (i.e., symmetric triazine) herbicide for the control of broad-leaf weeds. Approximately 800 million pounds were used in the United States between 1980 and 1990. As a result of this widespread use, for both selective and nonselective weed control, atrazine and other *s*-triazine-containing compounds have been detected in ground and surface water in several countries.

Numerous studies on the environmental fate of atrazine have shown that atrazine is a recalcitrant compound that is transformed to CO₂ very slowly, if at all, under aerobic or anaerobic conditions. It has a water solubility of 33 mg/l at 27°C. Its half-life (i.e., time required for half of the original concentration to dissipate) can vary from about 4 weeks to about 57 weeks when present at a low concentration (i.e., less than about 2 parts per million (ppm)) in soil. High concentrations of atrazine, such as those occurring in spill sites have been reported to dissipate even more slowly.

As a result of its widespread use, atrazine is often detected in ground water and soils in concentrations exceeding the maximum contaminant level (MCL) of 3 µg/l (i.e., 3 parts per billion (ppb)), a regulatory level that took effect in 1992. Point source spills of atrazine have resulted in levels as high as 25 ppb in some wells. Levels of up to 40,000 mg/l (i.e., 40,000 parts per million (ppm)) atrazine have been found in the soil at spill sites more than ten years after the spill incident. Such point source spills and subsequent runoff can cause crop damage and ground water contamination.

There have been numerous reports on the isolation of *s*-triazine-degrading microorganisms (see, e.g., Behki et al., J. Agric. Food Chem., **34**, 746-749 (1986); Behki et al., Appl. Environ. Microbiol., **59**, 1955-1959 (1993); Cook, FEMS Microbiol. Rev., **46**, 93-116 (1987); Cook et al., J. Agric. Food Chem., **29**, 1135-1143 (1981); Erickson et al., Critical Rev. Environ. Cont., **19**, 1-13 (1989); Giardina et al., Agric. Biol. Chem., **44**, 2067-2072 (1980); Jessee et al., Appl. Environ. Microbiol., **45**, 97-102 (1983); Mandelbaum et al., Appl. Environ. Microbiol., **61**, 1451-1457 (1995); Mandelbaum et al., Appl. Environ. Microbiol., **59**, 1695-1701 (1993); Mandelbaum et al., Environ. Sci. Technol.,

27, 1943-1946 (1993); Radosevich et al., Appl. Environ. Microbiol., 61, 297-302 (1995); and Yanze-Kontchou et al., Appl. Environ. Microbiol., 60, 4297-4302 (1994)). Many of the organisms described, however, failed to mineralize atrazine (see, e.g., Cook, FEMS Microbiol. Rev., 46, 93-116 (1987); and Cook et al., J. Agric. Food Chem., 29, 1135-1143 (1981)). While earlier studies have reported atrazine degradation only by mixed microbial consortia, more recent reports have indicated that several isolated bacterial strains can degrade atrazine. In fact, research groups have identified atrazine-degrading bacteria classified in different genera from several different locations in the U.S. (e.g., Minnesota, Iowa, Louisiana, and Ohio) and Switzerland (Basel).

An atrazine-degrading bacterial culture, identified as *Pseudomonas* sp. strain ADP (Mandelbaum et al., Appl. Environ. Microbiol., 61, 1451-1457 (1995); Mandelbaum et al., Appl. Environ. Microbiol., 59, 1695-1701 (1993); de Souza et al., J. Bact., 178, 4894-4900 (1996); and Mandelbaum et al., Environ. Sci. Technol., 27, 1943-1946 (1993)), was isolated and was found to degrade atrazine at concentrations greater than about 1,000 µg/ml under growth and non-growth conditions. See also, Radosevich et al., Appl. Environ. Microbiol., 61, 297-302 (1995) and Yanze-Kontchou et al., Appl. Environ. Microbiol., 60, 4297-4302 (1994). *Pseudomonas* sp. strain ADP (Atrazine Degrading *P*sedomonas) uses atrazine as a sole source of nitrogen for growth. The organism completely mineralizes the *s*-triazine ring of atrazine under aerobic growth conditions. That is, this bacteria is capable of degrading the *s*-triazine ring and mineralizing organic intermediates to inorganic compounds and ions (e.g., CO₂).

The genes that encode the enzymes for MELAMINE (2,4,6-triamino-*s*-triazine) metabolism have been isolated from a *Pseudomonas* sp. strain. The genes that encode atrazine degradation activity have been isolated from *Rhodococcus* sp. strains; however, the reaction results in the dealkylation of atrazine. In addition, the gene that encodes atrazine dechlorination has been isolated from a *Pseudomonas* sp. strain. See, for example, de Souza et al., Appl. Environ. Microbiol., 61, 3373 (1995). The protein expressed by the gene disclosed by de Souza et al., degrades atrazine, for example, at a V_{max} of about 2.6 µmol of hydroxyatrazine per min per mg protein. Although this is

significant, it is desirable to obtain genes and the proteins they express that are able to dechlorinate triazine-containing compounds with chlorine moieties at an even higher rate and/or under a variety of conditions, such as, but not limited to, conditions of high temperature (e.g., at least about 45°C and preferably at least about 65°C), various pH conditions, and/or under conditions of high salt content (e.g., about 20-30 g/L), or under other conditions in which the wild type enzyme is not stable, efficient, or active. Similarly, it is desirable to obtain genes and proteins encoded by these genes that degrade triazine-containing compounds such as those triazine containing compounds available under the trade names; "AMETRYN", "PROMETRYN", "CYANAZINE", "MELAMINE", "SIMAZINE", as well as TERBUTHYLAZINE and desethyldeisopropylatriazine. It is also desirable to identify proteins expressed in organisms that degrade triazine-containing compounds in the presence of other nitrogen sources such as ammonia and nitrate.

Summary of the Invention

The present invention provides isolated and purified DNA molecules that encode atrazine degrading enzymes similar to, but having different catalytic activities from a wild type (i.e., from an isolated but naturally occurring atrazine chlorohydrolase). The term "altered enzymatic activities" is used to refer to homologs of atrazine chlorohydrolase having altered catalytic rates as quantitated by k_{cat} and K_m , improved ability to degrade atrazine, altered substrate ranges, altered activities as compared to the native sequence in aqueous solutions, altered stability in solvents, altered active temperature ranges or altered reaction conditions such as salt concentration, pH, improved activity in a soil environment, and the like, as compared with the wild-type atrazine chlorohydrolase (AtzA) protein.

In one preferred embodiment, the present invention provides DNA fragments encoding a homolog of atrazine chlorhydrolase and comprising the sequence selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NOS:7-11 and SEQ ID NOS: 17-21. In one embodiment the invention relates to these DNA fragments in a vector, preferably an expression vector.

Further, the invention relates to the DNA fragment in a cell. In one embodiment the cell is a bacterium and in a preferred embodiment, the bacterium is *E. coli*.

The invention also relates to *s*-triazine-degrading proteins having at least one amino acid different from the protein of SEQ ID NO:2, wherein the
5 coding region of the nucleic acid encoding the *s*-triazine degrading protein has at least 95% homology to SEQ ID NO:1 and wherein the *s*-triazine-degrading protein has an altered catalytic activity as compared with the protein having the sequence of SEQ ID NO:2. In one embodiment, the protein is selected from the group consisting of SEQ ID NOS: 5, 6 and 22-26. In one embodiment the
10 substrate for the *s*-triazine degrading protein is ATRAZINE. In another embodiment the substrate for the *s*-triazine degrading protein is TERBUTHYLAZINE and in yet another embodiment the substrate for the *s*-triazine degrading protein is MELAMINE. In another embodiment this invention relates to a remediation composition comprising a cell producing at
15 least one *s*-triazine-degrading protein having at least one amino acid different from the protein of SEQ ID NO:2, wherein the coding region of the nucleic acid encoding the *s*-triazine degrading protein has at least 95% homology to SEQ ID NO:1 and wherein the *s*-triazine-degrading protein has an altered catalytic activity as compared with the protein having the sequence of SEQ ID NO:2. In a
20 preferred embodiment the composition is suitable for treating soil or water. In another embodiment the remediation composition comprises at least one *s*-triazine-degrading protein having at least one amino acid different from the protein of SEQ ID NO:2, wherein the coding region of the nucleic acid encoding the *s*-triazine degrading protein has at least 95% homology to SEQ ID NO:1 and
25 wherein the *s*-triazine-degrading protein has an altered catalytic activity as compared with the protein having the sequence of SEQ ID NO:2. In a preferred embodiment this composition is also suitable for treating soil or water. In one embodiment the remediation composition comprises the protein bound to an immobilization support. In yet another embodiment, these proteins are
30 homotetramers, such as the homotetramers formed by AtzA.

In another embodiment the invention relates to a protein selected from the group consisting of proteins comprising the amino acid sequences of SEQ ID NOS: 5, 6 and 22-26.

5 In another aspect of this invention, the invention relates to a DNA fragment having a portion of its nucleic acid sequence having at least 95% homology to a nucleic acid sequence consisting of position 236 and ending at position 1655 of SEQ ID NO:1, wherein the DNA fragment is capable of hybridizing under stringent conditions to SEQ ID NO:1 and wherein there is at least one amino acid change in the protein encoded by the DNA fragment as compared with SEQ ID NO:2 and wherein the protein encoded by the DNA fragment is capable of dechlorinating at least one *s*-triazine-containing compound and has a catalytic activity different from the enzymatic activity of the protein of SEQ ID NO:2. In one embodiment the *s*-triazine-containing compound is ATRAZINE, TERBUTHYLAZINE, or MELAMINE. In one
10 embodiment.

The invention also relates to a method for treating a sample comprising an *s*-triazine containing compound comprising the step of adding a adding a protein to a sample comprising an *s*-triazine-containing compound wherein the protein is encoded by gene having at least a portion of the nucleic acid sequence of the gene having at least 95% homology to the sequence beginning at position 236 and ending at position 1655 of SEQ ID NO:1, wherein the gene is capable of hybridizing under stringent conditions to SEQ ID NO:1, wherein there is at least one amino acid change in the protein encoded by the DNA fragment as compared with SEQ ID NO:2 and wherein the protein has an altered catalytic activity as compared to the protein having the amino acid sequence of SEQ ID NO:2. In one embodiment, the composition comprises bacteria expressing the protein. In one embodiment the *s*-triazine -containing compound is atrazine, in another the *s*-triazine-containing compound is TERBUTHYLAZINE and in another the *s*-triazine containing compound is
25 (2,4,6-triamino-*s*-triazine). In one embodiment, the protein encoded by the gene
30 is selected from the group consisting of SEQ ID NOS: 5, 6 and 22-26.

In another aspect, this invention relates to a method for obtaining homologs of an atrazine chlorohydrolase comprising the steps of obtaining a nucleic acid sequence encoding atrazine chlorohydrolase, mutagenizing the nucleic acid to obtain a modified nucleic acid sequence that encodes for a protein
5 having an amino acid sequence with at least one amino acid change relative to the amino acid sequence of the atrazine chlorohydrolase, screening the proteins encoded by the modified nucleic acid sequence; and selecting proteins with altered catalytic activity as compared to the catalytic activity of the atrazine chlorohydrolase. Preferably, the atrazine chlorohydrolase nucleic acid sequence
10 is SEQ ID NO:1. In one embodiment the altered catalytic activity is an improved ability to degrade ATRAZINE. In another embodiment, the altered catalytic activity is an altered substrate activity.

Other homologs with an improved rate of catalytic activity for atrazine include clones A40, A42, A44, A46 and A60 having nucleic acid
15 sequences (SEQ ID NOS:17-21, respectively). Other homologs capable of better degrading TERBUTHYLAZINE include A42, A44, A46 and A60 as well as A11 and A13.

20

Brief Description of the Drawings

Fig. 1. Nucleotide sequence alignment of wild type *atzA* (bottom sequence) from *Pseudomonas sp.* strain ADP and clone (A7) (SEQ ID NO:1 and SEQ ID NO:3).

25 **Fig. 2.** Nucleotide sequence alignment of wild type *atzA* (bottom sequence) from *Pseudomonas sp.* strain ADP and clone (T7) (SEQ ID NO: 1 and SEQ ID NO:4).

Fig. 3. Amino acid sequence alignment of wild type AtzA (bottom sequence) from *Pseudomonas sp.* strain ADP and clone (A7) (SEQ ID NO:2 and SEQ ID NO:5).

30

Fig. 4. Amino acid sequence alignment of wild type AtzA from *Pseudomonas sp.* strain ADP and clone (T7) (SEQ ID NO:2 and SEQ ID NO:6).

Fig. 5. Nucleotide sequence alignment of wild type *atzA* (SEQ ID NO:1, bottom sequence) from *Pseudomonas sp.* strain ADP and clone (A11).

Fig. 5(a) provides the sequence from nucleic acids 11-543 (SEQ ID NO:7), Fig. 5(b) provides the sequence from nucleic acids 454-901 (SEQ ID NO:8), Fig. 5(c) provides the sequence from 1458-1851 (SEQ ID NO:9; N in this sequence indicates that this nucleotide has not been verified) and Fig. 5(d) provides the sequence from nucleic acids 1125-1482 (SEQ ID NO:10) of clone A11. The "N" in these sequences refer to nucleic acids that are being verified.

Fig. 6. Nucleotide sequence alignment of a portion of the nucleic acid sequence of wild type *atzA* from *Pseudomonas sp.* strain ADP and nucleic acids 436-963 of clone (A13) (SEQ ID NO:11 and SEQ ID NO:1).

Fig. 7. is a histogram illustrating the TERBUTHYLAZINE degradative ability of two homologs of this invention (T7= sample 3 and A7 = sample 4). Fig. 7(a) illustrates the % of TERBUTHYLAZINE remaining after exposure to AtzA or a homolog. Fig. 7(b) illustrates the relative amount of hydroxyterbuthylazine as a measure of TERBUTHYLAZINE degradation.


Fig. 8. is another set of histograms illustrating the terbuthylazine degradative ability of three homologs A7, All, and T7. Figure 8(a) provides the % of TERBUTHYLAZINE remaining after a 15 minute exposure to the homolog in the presence or absence of the metals and additives of Samples 1-10. Figure 8(b) provides the relative amount of hydroxterbuthylazine in the presence or absence of the metals and compounds of Samples 1-10.

Fig. 9. is a comparison of PCR amplified fragments using two primers of the atrazine hydrochlorase gene from 6 different types of bacteria; *Pseudomonas sp.* strain ADP; *Ralstonia* strain M91-3; *Clavibacter (Clav.)*; *Agrobacterium* strain J14(a); ND (an organism with no genus assigned) strain 38/38; and *Alcaligenes* strain SG1 (SEQ ID NOS: 12-16).

Detailed Description of the Invention

The present invention provides isolated and purified DNA molecules, and isolated and purified proteins, involved in the degradation of s-triazine-containing compounds. The proteins encoded by the genes of this

invention are involved in the dechlorination and/or the deamination of *s*-triazine-containing compounds. The wild type AtzA protein can catalyze the dechlorination of *s*-triazine-containing compounds but not the deamination of these compounds. The dechlorination reaction occurs on *s*-triazine containing compounds that include a chlorine atom and at least one alkylamino side chain. Such compounds have the following general formula:



wherein $R^1 = \text{Cl}$, $R^2 = \text{NR}^4\text{R}^5$ (wherein R^4 and R^5 are each independently H or a C_{1-3} alkyl group), and $R^3 = \text{NR}^6\text{R}^7$ (wherein R^6 and R^7 are each independently H or a C_{1-3} alkyl group), with the proviso that at least one of R^2 or R^3 is an alkylamino group. As used herein, an "alkylamino" group refers to an amine side chain with one or two alkyl groups attached to the nitrogen atom. Examples of such compounds include atrazine (2-chloro-4-ethylamino-6-isopropylamino-1,3,5-*s*-triazine), desethylatrazine (2-chloro-4-amino-6-isopropylamino-*s*-triazine), desisopropylatrazine (2-chloro-4-ethylamino-6-amino-*s*-triazine), and SIMAZINE (2-chloro-4,6-diethylamino-*s*-triazine).

Triazine degradation activity is encoded by a gene that is localized to a 21.5-kb *EcoRI* fragment, and more specifically to a 1.9-kb *AvaI* fragment, of the genome of *Pseudomonas* sp. ADP (ADP is strain designation for Atrazine-degrading *Pseudomonas*) bacterium. Specifically, these genomic fragments encode proteins involved in *s*-triazine dechlorination. The rate of degradation of atrazine that results from the expression of these fragments in *E. coli* is comparable to that seen for native *Pseudomonas* sp. strain ADP; however, in contrast to what is seen with native *Pseudomonas* sp. strain ADP, this degradation in *E. coli* is unaffected by the presence of inorganic nitrogen sources like ammonium chloride. This is particularly advantageous for regions

contaminated with nitrogen-containing fertilizers or herbicides, for example. The expression of atrazine degradation activity in the presence of inorganic nitrogen compounds broadens the potential use of recombinant organisms for biodegradation of atrazine in soil and water.

5 Hydroxyatrazine formation in the environment was previously thought to result solely from the chemical hydrolysis of atrazine (Armstrong et al., Environ. Sci. Technol., 2, 683-689 (1968); deBruijn et al., Gene, 27, 131-149 (1984); and Nair et al., Environ. Sci. Technol., 26, 1627-1634 (1992)). Previous reports suggest that the first step in atrazine degradation by environmental
10 bacteria is dealkylation. Dealkylation produces a product that retains the chloride moiety and is likely to retain its toxicity in the environment. In contrast to these reports, AtzA dechlorinates atrazine and produces a detoxified product in a one-step detoxification reaction that is amenable to exploitation in the remediation industry. There remains a need for atrazine-degrading enzymes with
15 improved activity.

 As used herein, the gene encoding a protein capable of dechlorinating atrazine and originally identified in *Pseudomonas* sp. strain ADP and expressed in *E. coli* is referred to as "*atzA*", whereas the protein that it encodes is referred to as "AtzA." Examples of the cloned wild type gene
20 sequence and the amino acid sequence derived from the gene sequence are provided as SEQ ID NO:1 and SEQ ID NO:2 respectively. As also used herein, the terms atrazine chlorohydrolase (AtzA) protein, atrazine chlorohydrolase enzyme, or simply atrazine chlorohydrolase, are used interchangeably, and refer to an atrazine chlorohydrolase enzyme involved in the degradation of atrazine
25 and similar molecules as discussed above.

 A "homolog" of atrazine chlorohydrolase is an enzyme derived from the gene sequence encoding atrazine chlorohydrolase where the protein sequence encoded by the gene is modified by amino acid deletion, addition, substitution, or truncation but that nonetheless is capable of dechlorinating or
30 deaminating *s*-triazine containing compounds. In addition, the homolog of atrazine chlorohydrolase (AtzA) has a nucleic acid sequence that is different

from the *atzA* sequence (SEQ ID NO:1) and produces a protein with modified biological properties or, as used herein, "altered enzymatic activities." These homologs include those with altered catalytic rates as quantitated by k_{cat} and K_m , altered substrate ranges, altered activities as compared to the native sequence in aqueous solutions, altered stability in solvents, altered active temperature ranges or altered reaction conditions such as salt concentration, pH, improved activity in a soil environment, and the like, as compared with the wild-type atrazine chlorohydrolase (AtzA) protein. Thus, provided that two molecules possess enzymatic activity to an *s*-triazine-containing substrate and one molecule has the gene sequence of *atzA* (SEQ ID NO:1), the other is considered a homolog of that sequence where 1) the gene sequence of the homolog differs from SEQ ID NO:1 such that there is at least one amino acid change in the protein encoded by SEQ ID NO:1 (i.e., SEQ ID NO:2); 2) the homolog has different enzymatic characteristics from the protein encoded by SEQ ID NO:1 such as, but not limited to, an altered substrate preference, altered rate of activity, or altered conditions for enzymatic activity such as temperature, pH, salt concentration or the like, as discussed *supra*; and 3) where the coding region of the nucleic acid sequence encoding the variant protein has at least 95% homology to SEQ ID NO:1.

As used herein, the terms "isolated and purified" refer to the isolation of a DNA molecule or protein from its natural cellular environment, and from association with other coding regions of the bacterial genome, so that it can be sequenced, replicated, and/or expressed. Preferably, the isolated and purified DNA molecules of the invention comprise a single coding region. Thus, the present DNA molecules are preferably those consisting of a DNA segment encoding a homolog of atrazine chlorohydrolase.

Using the nucleic acid encoding the wild-type *atzA* sequence and the amino acid sequence of the wild-type enzyme AtzA, similar atrazine degrading enzymes were identified in other bacteria. In fact, sequencing of the *atzA* gene in the other bacteria demonstrated a homology of at least 99% to the *atzA* sequence, suggesting little evolutionary drift (see SEQ ID NOS:12-16).

Homologs of the *atzA* gene could not be identified in the genomes of bacteria that did not metabolize atrazine. This information supports the theory that the *atzA* gene evolved to metabolize *s*-triazine-containing compounds.

The studies assessing the prevalence and homology of the *atzA* gene in a variety of bacterial genera also suggest that *atzA* is likely to be a relatively young, i.e. recently evolved gene. That the gene is recently evolved is supported by the attributes of the gene and the protein encoded by the gene. For example: (i) the gene has a limited *s*-triazine range that includes atrazine and the structurally analogous herbicide SIMAZINE, but does not act on all *s*-triazines; (ii) the gene has a high sequence homology to genes isolated from other bacteria that produce proteins with atrazine-degrading activity; (iii) is not organized with the *atzB* and *atzC* genes in a contiguous arrangement such as an operon; (iv) the gene lacks the type of coordinate genetic regulation seen, for example, in aromatic hydrocarbon biodegradative pathway genes; (v) the wild-type gene was isolated from a spill site containing high atrazine levels and (vi) it is suggested to have been environmentally undetectable until the last few years.

Genes involved in reactions common to most bacteria and mammals are more highly evolved and have attained catalytic proficiency closer to theoretical perfection. Genes that have evolved more recently have not had the evolutionary opportunity to maximize the level of catalytic efficiency that they could theoretically obtain. These enzymes are suboptimal. Suboptimal enzymes include enzymes that have a second order rate constant, k_{cat}/K_m , that is orders of magnitude below the diffusion-controlled limit of enzyme catalysis, $3 \times 10^8 \text{ M}^{-1}\text{s}^{-1}$. These enzymes have the potential to evolve higher k_{cat} , lower K_m , or both. Enzymes with higher k_{cat} , lower K_m , or both would appear to have selective advantage as a biodegradative enzyme because less enzyme with higher activity would serve the same metabolic need and conserve ATP expended in enzyme biosynthesis. Optimized enzymes have the further advantage of having an improved commercial value resulting from their improved efficiency or improved activity under a defined set of conditions.

Thus, the *atzA* gene is, potentially, an *s*-triazine compound-degrading progenitor with the potential for improvement and modification. AtzA is a candidate for studies to generate homologs with improved activity, i.e., enhanced rate, altered pH preference, salt concentration and the like. The k_{cat}/K_M for atrazine chlorohydrolase purified from *Pseudomonas* ADP is $5 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$, 3
5 orders of magnitude below the theoretical catalytic limit. That all of the *atzA* homologous genes from a survey of atrazine-degrading bacteria are so structurally and catalytically similar suggest that the *atzA* gene and the AtzA protein can be improved and will be improved naturally over time. Indeed, most
10 biodegradative enzymes are orders of magnitude below diffusion limiting enzyme rates and, under this hypothesis, are also candidates for gene and protein modifications.

In one embodiment of this invention, a method is disclosed for selecting or screening modified and improved *atzA* gene sequences that encode
15 protein with improved enzymatic activity, whether the activity is enzymatic rate, using atrazine as a substrate, as compared to the wild-type sequence, or improved activity under any of a variety of reaction conditions including, but not limited to, elevated temperature, salt concentration, altered substrate range, solvent conditions, pH ranges, tolerance or stability to a variety of environmental
20 conditions, or other reaction conditions that may be useful in bioremediation processes. The method preferably includes the steps of obtaining the wild-type *atzA* gene sequence, mutagenizing the gene sequence to obtain altered *atzA* sequences, selecting or screening for clones expressing altered AtzA activity and selecting gene sequences encoding AtzA protein with improved *s*-triazine-degrading activity.
25

As a first step for practicing the method of this invention, the wild-type *atzA* sequence (SEQ ID NO:1) is incorporated into a vector or into nucleic acid that is suitable for a particular mutagenesis procedure. The wild type *atzA* gene was first obtained as a 1.9-kb *AvaI* genomic fragment that encodes an
30 enzyme that transforms atrazine to hydroxyatrazine, termed atrazine chlorohydrolase. Methods for obtaining this fragment are disclosed by de Souza

et al. (Appl. Environ. Microb. 61:3373-3378, (1995)). The gene, *atzA*, has one large ORF (open reading frame) and produces a translation product of about 473 amino acids. A particularly constant portion of this gene appears to occur at position 236 and end at position 1655 of SEQ ID NO:1. The wild-type *atzA* gene from *Pseudomonas* strain ADP includes 1419 nucleotides and encodes a polypeptide of 473 amino acids with an estimated M_r of 52,421 and a pI of 6.6. The gene also includes a typical *Pseudomonas* ribosome binding site, beginning with GGAGA, located 11 bp upstream from the proposed start codon. A potential stop codon is located at position 1655.

The wild-type *atzA* sequence can be obtained from a variety of sources including a DNA library, containing either genomic or plasmid DNA, obtained from bacteria believed to possess the *atzA* DNA. Alternatively the original isolate identified as containing the *atzA* DNA is described in U.S. Pat. No. 5,508,193 and can be accessed as a deposit from the American Type Culture Collection (ATCC No. 55464 Rockville, Maryland). Libraries can be screened using oligonucleotide probes, for example, to identify the DNA corresponding to SEQ ID NO:1. SEQ ID NO:1 can also be obtained by PCR (polymerase chain reaction) using primers selected using SEQ ID NO:1 and the nucleic acid obtained from the *atzA*-containing organism (ATCC No. 55464) deposited with the American Type Culture Collection.

Screening DNA libraries or amplifying regions from prokaryotic DNA using synthetic oligonucleotides is a preferred method to obtain the wild-type sequence of this invention. The oligonucleotides should be of sufficient length and sufficiently nondegenerate to minimize false positives. In a preferred strategy, the actual nucleotide sequence(s) of the probe(s) is designed based on regions of the *atzA* DNA, preferably outside of the reading frame of the gene (the translated reading frame begins at position 236 and ends at position 1655 of SEQ ID NO:1) that have the least codon redundancy.

Cloning of the open reading frame encoding *atzA* into the appropriate replicable vectors allows expression of the gene product, the AtzA enzyme, and makes the coding region available for further genetic engineering.

The types of mutagenesis procedures that are capable of generating a variety of gene sequences based on a parent sequence, *atzA* or a previously mutagenized or altered sequence of *atzA*, are known in the art and each method has a preferred vector format. In general, the mutagenesis procedures selected is one that generates at least one modified *atzA* sequence and preferably a population of modified *atzA* gene sequences. Selecting or screening procedures are used to identify preferred modified enzymes (i.e., homologs) from the pool of modified sequences.

There are a number of methods in use for creating mutant proteins in a library format from a parent sequence. These include the polymerase chain reaction (Leung, D.W. et al. Technique 1:11-15, (1989)), Bartel, D.P. et al. Science 261:1411-1418 (1993)), cassette mutagenesis (Arkīn, A. et al. Proc. Natl. Acad. Sci. USA 89:7811-7815 (1992), Oliphant, A.R. et al., Gene 44:177-183 (1986), Hermes, J.D. et al., Proc. Natl. Acad. Sci. USA 87:696-700 (1990), Delgrave et al. Protein Engineering 6:327-331, (1993), Delgrave et al. Bio/Technology 11:1548-1552 (1993), and Goldman, ER et al., Bio/Technology 10:1557-1561 (1992)), as well as methods that exploit the standard polymerase chain reaction, including, but not limited to, DNA recombination during *in vitro* PCR (Meyerhans, A. et al., Nucl. Acids Res. 18:1687-1691 (1990), and Marton et al. Nucl. Acids Res. 19:2423-2426, 1991)), *in vivo* site specific recombination (Nissim et al. EMBO J. 13:692-698 (1994), Winter et al. Ann. Rev. Immunol. 12:433-55 (1994)), overlap extension and PCR (Hayashi et al. Biotechniques 17:310-315 (1994)), applied molecular evolution systems (Bock, L. C. et al., Nature 355:564-566 (1992), Scott, J. K. et al., Science 249: 386-390 (1990), Cwirla, S.E. et al. Proc. Natl. Acad. Sci. USA 87:6378-6382 (1990), McCafferty, J. et al. Nature 348:552-554 (1990)), DNA shuffling systems, including those reported by Stemmer et al. (Nature 370:389-391 (1994) and Proc. Natl. Acad. Sci. (USA) 91:10747-10751 (1994) and International Patent Application Publication Number WO 95/22625), and random *in vivo* recombination (see Caren et al. Bio/Technology 12: 433-55 (1994), Caloger et al. FEMS

Microbiology Lett. 97:41-44 (1992), International Patent Application Publication Numbers WO91/01087, to Galizzi and WO90/07576 to Radman, et al.).

Preferably, the method produces libraries with large numbers of mutant nucleic acid sequences that can be easily screened or selected without undue experimentation. Those skilled in the art will recognize that screening and/or selection methods are well documented in the art and those of ordinary skill in the art will be able to use the cited methods as well as other references similarly describing the afore-mentioned methods to produce pools of variant sequences. Preferred strategies include methods for screening for degradative activity of the *s*-triazine-containing compound on nutrient plates containing the homolog-encoding bacteria or by use of colormetric assays to detect the release of chlorine ions. Preferred selection assays include methods for selecting for homolog-containing bacterial growth on or in a *s*-triazine containing medium.

In a preferred method of this invention, gene shuffling, also termed recursive sequence recombination, is used to generate a pool of mutated sequences of the *atzA* gene. In this method the *atzA* gene, alone or in combination with the *atzB* gene, is amplified, such as by PCR, or, alternatively, multiple copies of the gene sequence (*atzA* and *atzB*) are isolated and purified. The gene sequence is cut into random fragments using enzymes known in the art, including DNAase I. The fragments are purified and the fragments are incubated with single or double-stranded oligonucleotides where the oligonucleotides comprise an area of identity and an area of heterology to the template gene or gene sequence. The resulting mixture is denatured and incubated with a polymerase to produce annealing of the single-stranded fragments at regions of identity between the single-stranded fragments. Strand elongation results in the formation of a mutagenized double-stranded polynucleotide. These steps are repeated at least once. In this gene shuffling technique, recombination occurs between substantially homologous, but non-identical, sequences of the *atzA* gene. In the studies provided in Example 2, the *atzB* gene was not gene-shuffled.

In the technique, published by Stemmer et al. (Nature, *supra*), the reassembled product is amplified by PCR and cloned into a vector. Clones containing the shuffled gene are next used in selection or screening assays. Example 2 discloses the use of a gene shuffling technique to generate pools of modified *atzA* sequences. The gene shuffling technique of Example 2 was modified based on the Stemmer et al. references. In this technique, an entire plasmid containing the *atzA* and *atzB* gene in a vector was treated with DNAase I and fragments between 500 and 2000 bp were gel purified. The fragments were assembled in a PCR reaction as provided in Example 2.

Once intact gene sequences are reassembled, they are incorporated into a vector suitable for expressing protein encoded by the reassembled nucleic acid, or as provided in Example 1, where the gene sequences are already in a vector, the vector can be incorporated directly into an organism suitable for replicating the vector. The vector containing the *atzA* gene is also preferably incorporated into a host suitable for expressing the *atzA* gene. The host, generally an *E. coli* species, is used in assays to screen or select for clones expressing the AtzA protein under defined conditions. The type of organism can be matched to the mutagenesis procedure and in Example 2, a preferred organism was the *E. coli* strain NM522.

The assays suitable for use in this invention can take any of a variety of forms for determining whether a particular protein produced by the organism containing the variant *atzA* sequences expresses an enzyme capable of dechlorinating or deaminating *s*-triazine compounds. Therefore, the types of assays that could be used in this invention include assays that monitor the degradation of *s*-triazine-containing compounds including ATRAZINE, SIMAZINE or MELAMINE using any of a variety of methods including, but not limited to, HPLC analysis to assess substrate degradation; monitoring clearing of precipitable *s*-triazine containing substrates, such as atrazine or TERBUTHYLAZINE, on solid media by bacteria containing the homologs of this invention; growth assays in media containing soluble substrate, monitoring the amount of chlorine released, as described by Bergman et al., Anal. Chem.,

29, 241-243 (1957) or the amount of nitrogen released; evaluating the derivitized product using gas chromatography and/or mass spectroscopy, solid agar plate assays with varied salt, pH substrate, solvent, or temperature conditions, colorimetric assays such as those provided by Epstein, J. ("Estimation of
5 Microquantitation of Cyanide", (1947) *Analytical Chemistry* 19(4):272-276) and Habig and Jakoby ("Assays for Differentiation of Glutathione s-transferases, *Methods in Enzymology* 77:398-405) as well as radiolabelled assays to assess, for example, the release of radiolabel as a result of enzymatic activity.

In a preferred assay, clones are tested for their ability to degrade s-
10 triazine-containing compounds such as atrazine, SIMAZINE, TERBUTHYLAZINE (2-chloro-4-(ethylamino)-6-(tertiary butyl-amino)-1,3,5-triazine), desethylatrazine, desisopropylatrazine, MELAMINE, and the like. In these assays, atrazine, or another insoluble s-triazine-containing substrate, is incorporated into a nutrient agar plate as the sole nitrogen source.

15 Concentrations of atrazine or other s-triazine-containing compounds can vary in the plate from about 300 µg/ml to at least about 1000 µg/ml and in a preferred embodiment about 500 µg/ml atrazine is used on the plate. Many s-triazines are relatively insoluble compounds in water and a suspension in an agar plate produces a cloudy appearance. Bacteria capable of metabolizing the insoluble s-
20 triazine-containing compounds produce a clearing on the cloudy agar plate. An exemplary assays is a modified assay disclosed by Mandelbaum et al. (Appl. Environ. Microbiol. 61:1451-1453, (1995)) and provided in Example 2. In these assays LB medium can be used with the atrazine because *E. coli* expressing AtzA homologs support atrazine-degrading activity in the presence of other
25 nitrogen sources. The assay demonstrates atrazine degradation by observing clearing zones surrounding clones expressing homologs of AtzA.

Clones are selected from the insoluble substrate assay based on their ability to produce, for example, a clearing in the substrate-containing plates. Similarly, assay conditions can be modified such as, but not limited to, salt, pH,
30 solvent, temperature, and the like, to select clones encoding AtzA homologs capable of degrading a substrate under a variety of test conditions. For example,

the pH of the assay can be altered to a pH range of about 5 to about 9. These assays would likely use isolated homolog protein to permit an accurate assessment of the effect of pH. The assay, or a modification of the assay, suitable for elevated temperatures (such as a soluble assay) can employ elevated
5 temperature ranges, for example, between about 50° to about 80°C. The assays can also be modified to include altered salt concentrations including conditions equivalent to salt concentrations of about 2% to at least about 5% and preferably less than about 10% NaCl.

Clones identified as having altered enzymatic activity as compared
10 with the native enzyme are further assessed to rule out if the apparent enhanced activity of the enzyme is the result of a faster or more efficient AtzA protein production or whether the effect observed is the result of an altered *atzA* gene sequence. For example, in Example 2, the *atzA* was expressed to a high level using pUC18 as a preferred method to rule out higher *in vivo* activity due to
15 increased expression.

Once triazine-degrading colonies are identified with the desired characteristics, the AtzA homologs are isolated for further analysis. Clones containing putative faster enzyme(s) can be picked, grown in liquid culture, and the protein homolog can be purified, for example, as described (de Souza et al., J. Bacteriology, 178:4894-4900 (1996)). The genes encoding the homologs can be
20 modified, as known in the art, for extracellular expression or the homologs can be purified from bacteria. An exemplary method for protein purification is provided in Example 4. In a preferred method, protein was collected from bacteria using ammonium sulfate precipitation and further purified by HPLC
25 (see for example, de Souza et al., App. Envir. Microbio. 61:3373-3378 (1995)).

Using these methods, a number of homologs were identified. Homologs can be identified using the assays discussed in association with this invention including the precipitable substrate assays on solid agar as described by Mandelbaum, et al. (*supra*). Homologs identified using the methods of
30 Example 2 were separately screened for atrazine-degrading activity, for enhanced TERBUTHYLAZINE-degrading activity and for activity against other

s-triazine-containing compounds. An assay for TERBUTHYLAZINE degrading activity is provided in Example 6. Two homologs were found to have at least a 10 fold higher activity and contained 8 different amino acids than the native AtzA protein (A7 and T7, see Figs. 1-4). A subsequent round of DNA shuffling
5 starting with the homolog gene sequence yielded further improvements in activity (A11 and A13 corresponding to nucleic acid SEQ ID NOS: 7-10 and SEQ ID NO:11 respectively). This enzyme and other AtzA homologs (clones A40, A42, A44, A46, A60 corresponding to nucleic acid SEQ ID NOS: 17-21 and to protein SEQ ID NOS: 22-26, respectively) represent catabolic enzymes
10 modified in their biological activity. Preferred homologs identified in initial studies include A7, T7, A11, A44, and A46.

Homologs were also identified with altered substrate activity. Both homologs T7 and A7 were able to degrade TERBUTHYLAZINE better than the wild-type enzyme. Other homologs capable of degrading TERBUTHYLAZINE
15 include A42, A44, A46 and A60.

Atrazine chlorohydrolase converts a herbicide to a non-toxic, non-herbicidal, more highly biodegradable compound and the kinetic improvement of the homologs has important implications for enzymatic environmental remediation of this widely used herbicide. Less protein is
20 required to dechlorinate the same amount of atrazine. Importantly, the protein can also be used for degradation of the s-triazine-compound TERBUTHYLAZINE.

This invention also relates to nucleic acid and protein sequences identified from the homologs of this invention. Peptide and nucleic acid
25 fragments of these sequences are also contemplated and those skilled in the art can readily prepare peptide fragments, oligonucleotides, probes and other nucleic acid fragments based on the sequences of this invention. The homologs of this invention include those with an activity different from the native atrazine chlorohydrolase (AtzA) protein. As noted *supra*, an activity that is different
30 from the native atrazine chlorohydrolase protein includes enzymatic activity that is improved or is capable of functioning under different conditions such as salt

concentration, temperature, altered substrate, or the like. Preferably, the DNA encoding the homologs hybridize to a DNA molecule complementary to the wild-type coding region of a DNA molecule encoding wild-type AtzA protein, such as the sequence provided in SEQ ID NO:1, under high to moderate
5 stringency hybridization conditions. The homologs preferably have a homology of at least 95% to SEQ ID NO:1. As used herein, "high stringency hybridization conditions" refers to, for example, hybridization conditions in buffer containing 0.25 M Na_2HPO_4 (pH 7.4), 7% sodium dodecyl sulfate (SDS), 1% bovine serum albumin (BSA), 1.0 mM ethylene diamine tetraacetic acid (EDTA, pH 8) at
10 65°C, followed by washing 3x with 0.1% SDS and 0.1x SSC (0.1x SSC contains 0.015 M sodium chloride and 0.0015 M trisodium citrate, pH 7.0) at 65°C.

A number of homologs have been identified using the methods of this invention. For example, SEQ ID NO:3 is the gene sequence of a homolog A7 of the *atzA* gene that shows enhanced atrazine degradation activity and,
15 surprisingly, also demonstrated enhanced TERBUTHYLAZINE degradation activity. TERBUTHYLAZINE degradation experiments are provided in Example 6. The amino acid sequence of the enzyme encoded by SEQ ID NO:3 identified as SEQ ID NO:5. SEQ ID NO: 4 is the gene sequence of the homolog T7 of the *atzA* gene that shows enhanced atrazine degradation activity and
20 enhanced TERBUTHYLAZINE degradation activity. A summary of the TERBUTHYLAZINE degradation activity for T7 and A7 is provided in Example 6. SEQ ID NO:6 provides the amino acid sequence of the homolog encoded by SEQ ID NO:4. Fig. 1 provides the nucleotide sequence alignment of wild type *atzA* from SEQ ID NO:1 with SEQ ID NO:3 and Fig.2 provides the
25 nucleotide sequence alignment of SEQ ID NO:1 with SEQ ID NO:4. Fig. 3 provides the amino acid sequence alignment of SEQ ID NO:2, the amino acid sequence of the protein encoded by SEQ ID NO:1, with SEQ ID NO:5 and Fig. 4 provides the amino acid sequence alignment of SEQ ID NO:2 with SEQ ID NO:6. A review of the sequences encoding A7 and T7 indicate that both
30 homologs have a total of 8 amino acid changes relative to native AtzA (SEQ ID NO:2). Seven amino acid changes are common to both A7 and T7. The nucleic

acid sequences of other homologs with altered activity include A40 (nucleic acid SEQ ID NO:17; amino acid sequence SEQ ID NO:22); A42 (nucleic acid SEQ ID NO:18; amino acid sequence SEQ ID NO:23); A44 (nucleic acid SEQ ID NO:19; amino acid sequence SEQ ID NO:24); A46 (nucleic acid SEQ ID NO:20; amino acid sequence SEQ ID NO:25); and A60 (nucleic acid SEQ ID NO:21; amino acid sequence SEQ ID NO:26).

Without intending to limit the scope of this invention, the success attributed to the identification of homologs of AtzA may be based on the recognition that this protein is not evolutionarily mature. Therefore, not all gene sequences are good candidates as the starting material for identifying a number of biological variants of a particular protein and similarly, not all enzymes are amenable to the order of magnitude of rate enhancement by directed evolution using DNA shuffling or other methods. Without intending to limit the scope of this invention, it is believed that some enzymes are already processing substrates at their theoretical rate limit. In these cases, catalysis is limited by the physical diffusion of the substrate onto the catalytic surface of the enzyme. Thus, changes in the enzyme would not likely improve the rate of catalysis. Examples of enzymes that operate at or near catalytic "perfection" are triosephosphate isomerase, fumarase, and crotonase (available from the GenBank database system). Even biodegradative enzymes that hydrolyze toxic substrates fall into this class. For example, the phosphotriesterase that hydrolyzes paraoxon operates near enough to the diffusion limit and suggests that it would not be a good candidate for mutagenic methods to improve the catalytic rate constant of the enzyme with its substrate (see Caldwell et al., Biochem. 30:7438-7444 (1991)).

The gene sequences of this invention can be incorporated into a variety of vectors. Preferably, the vector includes a region encoding a homolog of AtzA and the vector can also include other DNA segments operably linked to the coding sequence in an expression cassette, as required for expression of the homologs, such as a promoter region operably linked to the 5' end of the coding DNA sequence, a selectable marker gene, a reporter gene, and the like.

The present invention also provides recombinant cells expressing the homologs of this invention. For example, DNA that expresses the homologs of this invention can be expressed in a variety of bacterial strains including *E. coli* sp. strains and *Pseudomonas* sp. strains. Other organisms include, but are not limited to, *Rhizobium*, *Bacillus*, *Bradyrhizobium*, *Arthrobacter*, *Alcaligenes*, and other rhizosphere and nonrhizosphere soil microbe strains.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable hosts for vectors encoding *atzA* or its homologs. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe*, *Kluyveromyces* hosts such as, e.g., *K. lactis*, *K. fragilis*, *K. bulgaricus*, *K. thermotolerans*, and *K. marxianus*, *Pichia pastoris*, *Candida*, *Trichoderma reesia*, *Neurospora crassa*, and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans*.

Prokaryotic cells used to produce the homologs of this invention are cultured in suitable media, as described generally in Maniatis et al., Molecular Cloning: A Laboratory Manual; Cold Spring Harbor Press: Cold Spring Harbor, NY (1989). Any necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. In general the *E. coli* expressing the homologs of this invention are readily cultured in LB media (see Maniatis, *supra*). The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to those skilled in the art. Induction of cells to express the AtzA protein is accomplished using the procedures required by the particular expression system selected. The host cells referred to in this disclosure are generally cultured *in vitro*. Cells are harvested, and cell extracts are prepared, using standard laboratory protocols.

This invention also relates to isolated proteins that are the product of the gene sequences of this invention. The isolated proteins are protein

homologs of the wild-type atrazine chlorohydrolase enzyme despite their potential for altered substrate preference. The protein can be isolated in a variety of methods disclosed in the art and a preferred method for isolating the protein is provided in Examples 4 and 5 and in the publications of de Souza et al. (*supra*).

5 The wild-type AtzA protein acts on Atrazine, desethylatrazine, Desisopropylatrazine and SIMAZINE but did not degrade Desethyldeisopropylatrazine or MELAMINE and only poorly degraded TERBUTHYLAZINE. Homologs identified in this invention have a spectrum of substrate preferences identical to the wild-type AtzA protein and in addition, for
10 example, are able to degrade other substrates such as TERBUTHYLAZINE. That homologs were identified that were capable of degrading two different *s*-triazine-containing compounds suggests that the methods of this invention can be used on the wild-type progenitor *atzA* gene or on the homologs produced by this invention to produce even more useful proteins for environmental remediation of
15 *s*-triazine-containing compounds. Example 7 provides an assay for detecting degradation, including deamination, of a soluble *s*-triazine-containing compound.

 Various environmental remediation techniques are known that utilize high levels of proteins. Bacteria or other hosts expressing the homologs
20 of this invention can be added to a remediation mix or mixture in need of remediation to promote contaminate degradation. Alternatively, isolated AtzA homologs can be added. Proteins can be bound to immobilization supports, such as beads, particles, films, etc., made from latex, polymers, alginate, polyurethane, plastic, glass, polystyrene, and other natural and man-made support materials.
25 Such immobilized protein can be used in packed-bed columns for treating water effluents. The protein can be used to remediate liquid samples, such as contaminated water, or solids. The advantage of some of the homologs identified thus far indicate that the homologs demonstrate an ability to degrade more than one substrate and to degrade the substrate at a faster rate or under different
30 reaction conditions from the native enzyme.

 All references and publications cited herein are expressly incorporated by reference into this disclosure. The invention will be further

described by reference to the following detailed examples. Particular
embodiments of this invention will be discussed in detail and reference has been
made to possible variations within the scope of this invention. There are a
variety of alternative techniques and procedures available to those of skill in the
art which would similarly permit one to successfully perform the intended
invention that do not detract from the spirit and scope of this invention.

Example 1

Isolation of Wild-type *atzA* gene from *Pseudomonas* sp. strain ADP

Bacterial strains and growth conditions.

Pseudomonas sp. strain ADP (Mandelbaum et al., Appl. Environ. Microbiol., 59, 1695-1701 (1993)) was grown at 37°C on modified minimal salt buffer medium, containing 0.5% (wt/vol) sodium citrate dihydrate. The atrazine stock solution was prepared as described in Mandelbaum et al., Appl. Environ. Microbiol., 61, 1451-1457 (1995)). *Escherichia coli* DH5 α was grown in Luria-Bertani (LB) or M63 minimal medium, which are described in Maniatis et al., Molecular Cloning: A Laboratory Manual; Cold Spring Harbor Press: Cold Spring Harbor, NY (1989). Tetracycline (15 μ g/ml), kanamycin (20 μ g/ml), and chloramphenicol (30 μ g/ml) were added as required.

To construct the *Pseudomonas* sp. strain ADP genomic library, total genomic DNA was partially digested with *Eco*RI, ligated to the *Eco*RI-digested cosmid vector pLAFR3 DNA, and packaged *in vitro*. The completed genomic DNA library contained 2000 colonies.

To identify the atrazine degrading clones, the entire gene library was replica-plated onto LB medium containing 500 μ g/ml atrazine and 15 μ g/ml tetracycline. Fourteen colonies having clearing zones were identified. All fourteen clones degraded atrazine, as determined by HPLC analysis. Cosmid DNA isolated from the fourteen colonies contained cloned DNA fragments which were approximately 22 kb in length. The fourteen clones could be subdivided into six groups on the basis of restriction enzyme digestion analysis using *Eco*RI. All fourteen clones, however, contained the same 8.7 kb *Eco*RI

fragment. Thirteen of the colonies, in addition to degrading atrazine, also produced an opaque material that surrounded colonies growing on agar medium. Subsequent experiments indicated that the opaque material only was observed in *E. coli* clones which accumulated hydroxyatrazine. Thus, the cloudy material surrounding *E. coli* pMD2-pMD4 colonies was due to the deposition of hydroxyatrazine in the growth medium. The one colony that degraded atrazine without the deposition of the opaque material was selected for further analysis. The clone from this colony was designated pMD1.

Example 2 Mutagenesis Procedure

Gene Shuffling. Atz A and B genes were subcloned from pMD1 into pUC18. The two inserts were reduced in size to remove extraneous DNA. A 1.9 kb *Ava*I fragment containing *atzA* was end-filled and cloned into the end-filled *Ava*I site of pUC18. A 3.9 kb *Cla*I fragment containing *atzB* was end-filled and cloned into the *Hinc*II site of pUC18. The gene *atzA* was then excised from pUC18 with *Eco*RI and *Bam*HI, *AtzB* with *Bam*HI and *Hind*III, and the two inserts were co-ligated into pUC18 digested with *Eco*RI and *Hind*III. The result was a 5.8 kb insert containing *AtzA* and *AtzB* in pUC18 (total plasmid size 8.4 kb).

Recursive sequence recombination was performed by modifications of existing procedures (Stemmer, W., Proc. Natl. Acad. Sci. USA 91:10747-10751 (1994) and Stemmer, W. Nature 370:389-391 (1994)). [Mervyn, do you know more now about what was done?] The entire 8.4 kb plasmid was treated with DNAase I in 50 mM Tris-Cl pH 7.5, 10 mM $MnCl_2$ and fragments between 500 and 2000 bp were gel purified. The fragments were assembled in a PCR reaction using Tth-XL enzyme and buffer from Perkin Elmer, 2.5 mM $MgOAc$, 400 μ M dNTPs and serial dilutions of DNA fragments. The assembly reaction was performed in an MJ Research "DNA Engine" thermocycler programmed with the following cycles:

- 1 94°C, 20 seconds
- 2 94°C, 15 seconds
- 3 40°C, 30 seconds
- 4 72°C, 30 seconds + 2 seconds per cycle
- 5 go to step 2 39 more times
- 6 4°C

The *atzA* gene could not be amplified from the assembly reaction using the polymerase chain reaction, so instead DNA from the reaction was purified by standard phenol extraction and ethanol precipitation methods and digested with KpnI to linearize the plasmid (the KpnI site in pUC18 was lost during subcloning, leaving only the KpnI site in *atzA*). Linearized plasmid was gel-purified, self-ligated overnight and transformed into *E coli* strain NM522.

Serial dilutions of the transformation reaction were plated onto LB plates containing 50 µg/ml ampicillin, the remainder of the transformation was stored in 25% glycerol and frozen at -80°C. Once the transformed cells were titered, the frozen cells were plated at a density of between 200 and 500 on 150 mm diameter plates containing 500 µg/ml atrazine or another substrate and grown at 37°C.

Atrazine at 500 µg/ml forms an insoluble precipitate creating a cloudy appearance on the agar plate. The solubility of atrazine is about 30 µg/ml, therefore for precipitable substrate assays, such as the assay disclosed here, the atrazine concentration should be preferably greater than 30 µg/ml. Atrazine or hydroxyatrazine were incorporated in solid LB or minimal medium, as described in Mandelbaum et al., Appl. Environ. Microbiol., **61**, 1451-1457 (1995), at a final concentration of 500 µg/ml to produce an opaque suspension of small particles in the clear agar. AtzA and the homologs with atrazine-degrading activity convert atrazine into a soluble product. The degradation of atrazine or hydroxyatrazine by wild-type and recombinant bacteria was indicated by a zone of clearing surrounding colonies. The more active the homolog, the more rapidly a clear halo formed on atrazine-containing plates. Positive colonies that most rapidly formed the largest clear zones were selected initially for further analysis. The (approximately) 40 best colonies were picked, pooled, grown in the presence of 50 µg/ml ampicillin and plasmid prepared from them. More

efficient enzymes can also be tested using atrazine concentrations greater than 500 µg/ml.

The entire process (from DNAase-treatment to plating on atrazine plates) was repeated 4 times as a method for further improving on the rate of enzymatic activity. In several experiments, cells were plated on plates containing 500 µg/ml atrazine and on plates containing 500 µg/ml of the atrazine analogue TERBUTHYLAZINE.

Other compounds can be tested in similar assays replacing atrazine (2-chloro-4-ethylamino-6-isopropylamino-1,3,5-s-triazine) for the following compounds: desethylatrazine (2-chloro-4-amino-6-isopropylamino-s-triazine), deisopropylatrazine (2-chloro-4-ethylamino-6-amino-s-triazine), hydroxyatrazine (2-hydroxy-4-ethylamino-6-isopropylamino-s-triazine), desethylhydroxyatrazine (2-hydroxy-4-amino-6-isopropylamino-s-triazine), desisopropylhydroxyatrazine (2-hydroxy-4-amino-6-isopropylamino-s-triazine), desethyldeisopropylatrazine (2-chloro-4,6-diamino-s-triazine), SIMAZINE (2-chloro-4,6-diethylamino-s-triazine), TERBUTHYLAZINE (2-chloro-4-ethylamino-6-terbutylamino-s-triazine, and MELAMINE (2,4,6-triamino-s-triazine) were obtained from Ciba Geigy Corp., Greensboro, N.C. Ammelide (2,4-dihydroxy-6-amino-s-triazine), ammeline (2-hydroxy-4,6,-diamino-s-triazine) were obtained from Aldrich Chemical Co., Milwaukee, WI.

Example 3 DNA Sequencing of Wild-Type *atxA* and Homolog *atxA* genes

DNA Sequencing. The nucleotide sequence of the approximately 1.9-kb *Ava*I DNA fragment in vector pACYC184, designated pMD4, or the homologs in pUC18 or another vector was determined using both DNA strands. DNA was sequenced by using a PRISM Ready Reaction DyeDeoxy Terminator Cycle Sequencing kit (Perkin-Elmer Corp., Norwalk, CT) and a ABI Model 373A DNA Sequencer (Applied Biosystems, Foster City, CA). Nucleotide sequence was determined initially by subcloning and subsequently by using primers designed based on sequence information obtained from subcloned DNA fragments. The GCG sequence analysis software package (Genetics Computer

Group, Inc., Madison, WI) was used for all DNA and protein sequence comparisons. Radiolabelled chemicals were obtained from Ciba Geigy Corp., Greensboro, N.C.

5

Example 4 Protein Purification of AtzA or Homologs

E. coli transformed with a vector containing the wild type *atzA* gene or alternatively with a homolog, in a vector capable of directing expression of the gene as a protein, was grown overnight at 37°C in eight liters of LB medium containing 25 µg/ml chloramphenicol. The culture medium was centrifuged at 10,000 x g for 10 minutes at 4°C, washed in 0.85% NaCl, and the cell pellet was resuspended in 50 ml of 25 mM MOPS buffer (3-[N-morpholino]propane-sulfonic acid, pH 6.9), containing phenylmethylsulfonylfluoride (100 µg/ml). The cells were broken by three passages through an Amicon French Pressure Cell at 20,000 pounds per square inch (psi) at 4°C. Cell-free extract was obtained by centrifugation at 10,000 x g for 15 minutes. The supernatant was clarified by centrifugation at 18,000 x g for 60 minutes and solid NH₄SO₄ was added, with stirring, to a final concentration of 20% (wt/vol) at 4°C. The solution was stirred for 30 minutes at 4°C and centrifuged at 12,000 x g for 20 minutes. The precipitated material was resuspended in 50 ml of 25 mM MOPS buffer (pH 6.9), and dialyzed overnight at 4°C against 1 liter of 25 mM MOPS buffer (pH 6.9).

Where purified protein was desired, the solution was loaded onto a Mono Q HR 16/10 Column (Pharmacia LKB Biotechnology, Uppsala, Sweden). The column was washed with 25 mM MOPS buffer (pH 6.9), and the protein was eluted with a 0-0.5 M KCl gradient. Protein eluting from the column was monitored at 280 nm by using a Pharmacia U.V. protein detector. Pooled fractions containing the major peak were dialyzed overnight against 1 liter 25 mM MOPS buffer (pH 6.9). The dialyzed material was assayed for atrazine degradation ability by using HPLC analysis (see above) and analyzed for purity by sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoreses (Laemmli).

Protein Verification: Protein subunit sizes were determined by SDS polyacrylamide gel electrophoresis by comparison to known standard proteins, using a Mini-Protean II gel apparatus (Biorad, Hercules, CA). The size of the holoenzyme was determined by gel filtration chromatography on a
5 Superose 6 HR (1.0 x 30.0 cm) column, using an FPLC System (Pharmacia, Uppsala, Sweden). The protein was eluted with 25 mM MOPS buffer (pH 6.9) containing 0.1 M NaCl. Proteins with known molecular weights were used as chromatography standards. Isoelectric point determinations were done using a Pharmacia Phast-Gel System and Pharmacia IEF 3-9 media. A Pharmacia broad-
10 range pI calibration kit was used for standards.

Enzyme Kinetics. Purified AtzA protein and homologs of the protein at 50 µg/ml, were separately added to 500 µl of different concentrations of atrazine (23.3 µM, 43.0 µM, 93 µM, 233 µM, and 435 µM in 25 mM MOPS buffer, pH 6.9) or another *s*-triazine-containing compound and reactions were
15 allowed to proceed at room temperature for 2, 5, 7, and 10 minutes. The reactions were stopped by boiling the reaction tubes at specific times, the addition of 500 µl acetonitrile and rapid freezing at -80°C. Thawed samples were centrifuged at 14,000 rpm for 10 minutes, the supernatants were filtered through a 0.2 µM filter, and placed into crimp-seal HPLC vials. HPLC analysis was done
20 as described above. Based on HPLC data, initial rates of atrazine degradation and hydroxyatrazine formation were calculated and Michaelis Menton and Lineweaver Burke plots were constructed.

Effect of simple nitrogen sources on atrazine degradation.
From experiments done with *Pseudomonas* species strain ADP on solid media
25 with 500 ppm atrazine and varying concentrations of ammonium chloride, ammonium chloride concentrations as low as 0.6-1.2 mM were sufficient to inhibit visible clearing on the plates, even after 2 weeks of incubation either at 28°C or 37°C. With similar experiments using *E. coli* DH5α (pMD1 or pMD2) and other *E. coli* strains, atrazine degradation was observed in the presence of
30 ammonium chloride concentrations as high as 48 mM. This value is almost 40-80 fold higher than the wild-type tolerance for ammonium chloride with

concomitant atrazine degradation. Therefore, it was not necessary to use media free of exogenous ammonia in the screening assays.

Example 5

Further characterization of the enzymatic activity of the homologs

Analysis of atrazine metabolism by *E. coli* clones. The extent and rate of atrazine degradation was determined in liquid culture. *E. coli* clones containing plasmids capable of expressing the homologs were compared to *Pseudomonas* sp. strain ADP for their ability to transform ring-labelled [¹⁴C]-atrazine to water-soluble metabolites. This method, which measures [¹⁴C]-label partitioning between organic and aqueous phases, had previously been used with *Pseudomonas* sp. ADP to show the transformation of atrazine to metabolites that partition into the aqueous phase, in Mandelbaum et al., Appl. Environ. Microbiol., 61, 1451-1457 (1995). When *Pseudomonas* sp. strain ADP or *E. coli* capable of expressing the homologs of this invention were incubated for 2 hours with [¹⁴C]-atrazine, 98%, 97%, 88%, and 92%, respectively, of the total recoverable radioactivity was found in the aqueous phase. Greater than 90% of the initial radioactivity was accounted for as atrazine plus water soluble metabolites, indicating that little or no ¹⁴CO₂ was formed. In contrast, forty-four percent of the radioactivity was lost from the *Pseudomonas* ADP culture after 18.5 hours. In previous studies done with *Pseudomonas* sp. strain ADP and ring-labelled ¹⁴C-atrazine, radiolabel was lost from culture filtrates as ¹⁴CO₂ (see, e.g., Mandelbaum et al., Appl. Environ. Microbiol., 61, 1451-1457 (1995)). Retention of the radiolabel is indicative of lack or inhibition of enzymatic activity. While these studies were performed for AtzA, similar studies are used to assess the activity of the homologs of this invention.

Example 6

Assays to detect homologs of AtzA on TERBUTHYLAZINE

TERBUTHYLAZINE was incorporated in solid LB medium at a final concentration of about 400-500 µg/ml to produce an opaque suspension of sample particles in the clear agar. The degradation of terbuthylazine by

recombinant bacteria was indicated by a zone of clearing surrounding the colonies. HPLC analysis was performed with a Hewlett Packard HP 1090 Liquid Chromatograph system equipped with a photodiode array detector and interfaced to an HP 79994A Chemstation. TERBUTHYLAZINE and its
5 metabolites were resolved by using an analytical C¹⁸ reverse-phase Nova-Pak HPLC column (4- μ m-diameter spherical packing, 150 by 3.9 mm; Waters Chromatography, Milford, Mass.) and an acetonitrile (ACN) gradient, in water, at a flow rate of 1.0 ml min⁻¹. Linear gradients of 0 to 6 min, 10 to 25% ACN; 6 to 21 min, 25 to 65% ACN; 21 to 23 min, 65 to 100% ACN; and 23 to 25 min,
10 100% ACN were used. Spectral data of the column eluent were acquired between 200 and 400 nm (12-nm bandwidth per channel) at a sampling frequency of 640 ms. Spectra were referenced against a signal of 500 nm.

Comparative results of an assay to assess TERBUTHYLAZINE degradation is provided in Figures 7 and 8. Figure 7 (a) provides a histogram
15 demonstrating the relative percentage of TERBUTHYLAZINE remaining in samples tested while Figure 7(b) provides a measure of the production of hydroxyterbuthylazine as a measure of TERBUTHYLAZINE degradation. Sample 1 is a control sample without enzyme. Sample 2 uses a two fold excess of AtzA protein as compared to the concentration of homolog added in Sample 3
20 and Sample 4. Sample 3 employed the T7 homolog (SEQ ID NO:6) and Sample 4 employed the A7 homolog (SEQ ID NO:5). Results were determined by HPLC as described above. Figure 8(a) provides the percentage of TERBUTHYLAZINE remaining after a 15 minute exposure to homologs A7, A11, and T7. Samples 1-10 refer to the effect of homolog activity in the
25 presence of 50 μ M of: Manganese (1); Manganese (2), EDTA (3); cobalt (4); zinc (5); iron (6); copper (7); nickel (8); no metal (9); or no enzyme (10). Figure 8(b) provides the relative amount of hydroxyterbuthylazine as a measure of TERBUTHYLAZINE degradation for homologs A7 (solid bar), A11 (hatched bar), or T7 (open bar) in the presence or absence of additives 1-10 (*supra*).

Example 7

Assays to detect homologs of AtzA on "MELAMINE"

5 "MELAMINE" (2, 4, 6-triamino -s-triazine) at a concentration of at least about 1 mM to about 5 mM and preferably about 2 mM MELAMINE is incorporated into solid minimal nutrient media as the sole nitrogen source. Bacteria are distributed on the plate and growth of the organisms is indicative of their ability to degrade MELAMINE, thereby releasing ammonia for growth.

10 Growth is evidence of the ability of the organisms expressing the homologs of this invention to deaminate MELAMINE. There is more than one nitrogen-containing group in MELAMINE. Therefore the selection of larger colonies on MELAMINE containing solid minimal nutrient media could be used to select for faster MELAMINE-degrading homologs.

15 A comparison of the nucleic acid sequence from a wild type MELAMINE degrading *Pseudomonas* NRRLB 12227 strain as compared to the *atzA* gene sequence indicated a homology of more than 90% over a 500 base pair sequence obtained from NRRLB using primer selected that were internal to *atzA* suggesting that homologs of *atzA* could be identified that degrade

20 "MELAMINE." This strain did not degrade atrazine. Moreover, homologs identified using the methods of Example 2 are subjected to further mutagenesis and colonies capable of growing in MELAMINE can be identified. Colonies containing the protein AtzA are tested for growth in MELAMINE under identical conditions. Other s-triazine containing compounds such as the

25 pesticides available under the tradenames "AMETRYN", "PROMETRYN", "PROMETRON", "ATRATON" and "CYROMAZINE" could also function as substrates for other homologs of this invention.

30 It will be appreciated by those skilled in the art that while the invention has been described above in connection with particular embodiments and examples, the invention is not necessarily so limited and that numerous other embodiments, examples, uses, modifications and departures from the

embodiments, examples and uses may be made without departing from the inventive scope of this application.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: REGENTS OF THE UNIVERSITY OF MINNESOTA
- (ii) TITLE OF INVENTION: DNA MOLECULES AND PROTEIN DISPLAYING
IMPROVED TRIAZINE COMPOUND DEGRADING ABILITY
- (iii) NUMBER OF SEQUENCES: 26
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: MUETING, RAASCH & GEBHARDT, P.A.
 - (B) STREET: 119 North Fourth Street
 - (C) CITY: Minneapolis
 - (D) STATE: Minnesota
 - (E) COUNTRY: USA
 - (F) ZIP: 55401
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) PRIORITY APPLICATION DATA:
 - (A) APPLICATION NUMBER: 60/035,404
 - (B) FILING DATE: 17-JAN-1997
 - (C) CLASSIFICATION:
- (vii) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: Not Assigned
 - (B) FILING DATE: 16-JAN-1998
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: MCCORMACK, MYRA M.
 - (B) REGISTRATION NUMBER: 36,602
 - (C) REFERENCE/DOCKET NUMBER: 110.00400201
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 612-305-1225
 - (B) TELEFAX: 612-305-1228

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1858 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTCGGGTAAC TTCTTGAGCG CGGCCACAGC AGCCTTGATC ATGAAGGCGA GCATGGTGAC	60
CTTGACGCCG CTCTTTTCGT TCTCTTTGTT GAACTGCACG CGAAAGGCTT CCAGGTCGGT	120
GATGTCCGCG TCGTCGTGGT TGGTGACGTG CGGGATGACC ACCCAGTTGC GGTGCAGGTT	180
TTTCGATGGC ATAATATCTG CGTTGCGACG TGTAACACAC TATTGGAGAC ATATCATGCA	240
AACGCTCAGC ATCCAGCACG GTACCCTCGT CACGATGGAT CAGTACCGCA GAGTCCTTGG	300
GGATAGCTGG GTTCACGTGC AGGATGGACG GATCGTCGCG CTCGGAGTGC ACGCCGAGTC	360
GGTGCTCCG CCAGCGGATC GGGTGATCGA TGCACGCGGC AAGGTCGTGT TACCCGGTTT	420
CATCAATGCC CACACCCATG TGAACCAGAT CCTCCTGCGC GGAGGGCCCT CGCACGGACG	480
TCAATTCTAT GACTGGCTGT TCAACGTTGT GTATCCGGGA CAAAAGGCGA TGAGACCGGA	540
GGACGTAGCG GTGGCGGTGA GGTGTATTG TGCGGAAGCT GTGCCGACG GGATTACGAC	600
GATCAACGAA AACGCCGATT CGGCCATCTA CCCAGGCAAC ATCGAGGCCG CGATGGCGGT	660
CTATGGTGAG GTGGGTGTGA GGGTCGTCTA CGCCCGCATG TTCTTTGATC GGATGGACGG	720
GCGCATTCAA GGGTATGTGG ACGCCTTGAA GGCTCGCTCT CCCCAGTCG AACTGTGCTC	780
GATCATGGAG GAAACGGCTG TGGCCAAAGA TCGGATCACA GCCCTGTCAG ATCAGTATCA	840
TGGCACGGCA GGAGGTCGTA TATCAGTTTG GCCCGCTCCT GCCACTACCA CGGCGGTGAC	900
AGTTGAAGGA ATGCGATGGG CACAAGCCTT CGCCCGTGAT CGGGCGGTAA TGTGGACGCT	960
TCACATGGCG GAGAGCGATC ATGATGAGCG GATTCATGGG ATGAGTCCCG CCGAGTACAT	1020
GGAGTGTTAC GGA CTCTTG ATGAGCGTCT GCAGGTCGCG CATTGCGTGT ACTTTGACCG	1080
GAAGGATGTT CGGCTGCTGC ACCGCCACAA TGTGAAGGTC GCGTCGCAGG TTGTGAGCAA	1140
TGCCTACCTC GGCTCAGGGG TGGCCCCCGT GCCAGAGATG GTGGAGCGCG GCATGGCCGT	1200
GGGCATTGGA ACAGATAACG GGAATAGTAA TGACTCCGCA AACATGATCG GAGACATGAA	1260
GT TTATGGCC CATATTCACC GCGCGGTGCA TCGGGATGCG GACGTGCTGA CCCCAGAGAA	1320

GATTCTTGAA ATGGCGACGA TCGATGGGGC GCGTTCGTTG GGAATGGACC ACGAGATTGG 1380
 TTCCATCGAA ACCGGCAAGC GCGCGGACCT TATCCTGCTT GACCTGCGTC ACCTCAGACG 1440
 ACTCTCACAT CATTGGCGG CCACGATCGT GTTTCAGGCT TACGGCAATG AGGTGGACAC 1500
 TGTCTGATT GACGGAAACG TTGTGATGGA GAACCGCCGC TTGAGCTTTC TTCCCCCTGA 1560
 ACGTGAGTTG GCGTTCCTTG AGGAAGCGCA GAGCCGCGCC ACAGCTATTT TGCAGCGGGC 1620
 GAACATGGTG GCTAACCCAG CTTGGCGCAG CCTCTAGGAA ATGACGCCGT TGCTGCATCC 1680
 GCCGCCCCCTT GAGGAAATCG CTGCCATCTT GGCGCGGCTC GGATTGGGGG GCGGACATGA 1740
 CCTTGATGGA TACAGAATTG CCATGAATGC GGCATTCCG TCCTTCGCTC GTGTGGAATC 1800
 GTTGGTAGGT GAGGGTCGAC TCGGGGCGCC AGCTTCCCGA AGAGGTGAAA GGCCCCGAG 1858

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 473 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gln Thr Leu Ser Ile Gln His Gly Thr Leu Val Thr Met Asp Gln
 1 5 10 15
 Tyr Arg Arg Val Leu Gly Asp Ser Trp Val His Val Gln Asp Gly Arg
 20 25 30
 Ile Val Ala Leu Gly Val His Ala Glu Ser Val Pro Pro Pro Ala Asp
 35 40 45
 Arg Val Ile Asp Ala Arg Gly Lys Val Val Leu Pro Gly Phe Ile Asn
 50 55 60
 Ala His Thr His Val Asn Gln Ile Leu Leu Arg Gly Gly Pro Ser His
 65 70 75 80
 Gly Arg Gln Phe Tyr Asp Trp Leu Phe Asn Val Val Tyr Pro Gly Gln
 85 90 95
 Lys Ala Met Arg Pro Glu Asp Val Ala Val Ala Val Arg Leu Tyr Cys
 100 105 110
 Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn Ala Asp
 115 120 125

Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val Tyr Gly
 130 135 140

Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp Arg Met
 145 150 155 160

Asp Gly Arg Ile Gln Gly Tyr Val Asp Ala Leu Lys Ala Arg Ser Pro
 165 170 175

Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val Ala Lys Asp
 180 185 190

Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly Gly Arg
 195 200 205

Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr Val Glu
 210 215 220

Gly Met Arg Trp Ala Gln Ala Phe Ala Arg Asp Arg Ala Val Met Trp
 225 230 235 240

Thr Leu His Met Ala Glu Ser Asp His Asp Glu Arg Ile His Gly Met
 245 250 255

Ser Pro Ala Glu Tyr Met Glu Cys Tyr Gly Leu Leu Asp Glu Arg Leu
 260 265 270

Gln Val Ala His Cys Val Tyr Phe Asp Arg Lys Asp Val Arg Leu Leu
 275 280 285

His Arg His Asn Val Lys Val Ala Ser Gln Val Val Ser Asn Ala Tyr
 290 295 300

Leu Gly Ser Gly Val Ala Pro Val Pro Glu Met Val Glu Arg Gly Met
 305 310 315 320

Ala Val Gly Ile Gly Thr Asp Asn Gly Asn Ser Asn Asp Ser Ala Asn
 325 330 335

Met Ile Gly Asp Met Lys Phe Met Ala His Ile His Arg Ala Val His
 340 345 350

Arg Asp Ala Asp Val Leu Thr Pro Glu Lys Ile Leu Glu Met Ala Thr
 355 360 365

Ile Asp Gly Ala Arg Ser Leu Gly Met Asp His Glu Ile Gly Ser Ile
 370 375 380

Glu Thr Gly Lys Arg Ala Asp Leu Ile Leu Leu Asp Leu Arg His Leu
 385 390 395 400

Arg Arg Leu Ser His His Leu Ala Ala Thr Ile Val Phe Gln Ala Tyr
 405 410 415

Gly Asn Glu Val Asp Thr Val Leu Ile Asp Gly Asn Val Val Met Glu
 420 425 430

Asn Arg Arg Leu Ser Phe Leu Pro Pro Glu Arg Glu Leu Ala Phe Leu
435 440 445

Glu Glu Ala Gln Ser Arg Ala Thr Ala Ile Leu Gln Arg Ala Asn Met
450 455 460

Val Ala Asn Pro Ala Trp Arg Ser Leu
465 470

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1808 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3;

GCGAGCATGG	TGACCTTGAC	GCCGCTCTTT	TCGTTCTCTT	TGTTGAACTG	CACGCGAAAG	60
GCTTCCAGGT	CGGTGATGTC	CGCGTCGTCG	TGGTTGGTGA	CGTGCGGGAT	GACCACCCAG	120
TTGCGGTGCA	GGTTTTTCGA	TGGCATAATA	TCTGCGTTGC	GACGTGTAAC	AACTATTGG	180
AGACATATCA	TGCAAACGCT	CAGCATCCAG	CACGGTACCC	TCGTCACGAT	GGATCAGTAC	240
CGCAGAGTCC	TTGGGGATAG	CTGGGTTTAC	GTGCAGGATG	GACGGATCGT	CGCGCTCGGA	300
GTGCACGCCG	AGTCGGTGCC	TCCGCCAGCG	GATCGGGTGA	TCGATGCACG	CGGCAAGGTC	360
GTGTTACCCG	GTTTCATCAA	TGCCCACACC	CATGTGAACC	AGATCCTCCT	GCGCGGAGGG	420
CCCTCGCACG	GGCGTCAATT	CTATGACTGG	CTGTTCAACG	TTGTGTATCC	GGGACAAAAG	480
GCGATGAGAC	CGGAGGACGT	AGCGGTGGCG	GTGAGGTTGT	ATTGTGCGGA	AGCTGTGCGC	540
AGCGGGATTA	CGACGATCAA	CGAAAACGCC	GATTCGGCCA	TCTACCCAGG	CAACATCGAG	600
GCCGCGATGG	CGGTCTATGG	TGAGGTGGGT	GTGAGGGTCG	TCTACGCCCC	CATGTTCTTT	660
GATCGGATGG	ACGGGCGCAT	TCAAGGGTAT	GTGGACGCCT	TGAAGGCTCG	CTCTCCCCAA	720
GTCGAACTGT	GCTCGATCAT	GGAGGGAACG	GCTGTGGCCA	AAGATCGGAT	CACAGCCCTG	780
TCAGATCAGT	ATCATGGCAC	GGCAGGAGGT	CGTATATCAG	TTTGGCCCCG	TCCTGCCACT	840
ACCACGGCGG	TGACAGTTGA	AGGAATGCGA	TGGGCACAAG	CCTTCGCCCC	TGATCGGGCG	900
GTAATGTGGA	CGCTTCACAT	GGCGGAGAGC	GATCATGATG	AGCGGATTCA	TGGGATGAGT	960

CCCCCGAGT ACATGGAGTG TTACGGACTC TTGGATGAGC GTCTGCAGGT CGCGCATTGC	1020
GTGTACTTTG ACCGGAAGGA TGTTCGGCTG CTGCACCGCC ACAATGTGAA GGTCGCGTCG	1080
CAGGTTGTGA GCAATGCCTA CCTCGGCTCA GGGGTGGCCC CCGTGCCAGA GATGGTGGAG	1140
CGCGGCATGG CCGTGGGCAT TGAACAGAT AACGGGAATA GTAATGACTC CGTAAACATG	1200
ATCGGAGACA TGAAGTTTAT GGCCCATATT CACCGCGCGG TGCATCGGGA TGCGGACGTG	1260
CTGACCCCAG AGAAGATTCT TGAAATGGCG ACGATCGATG GGGCGCGTTC GTTGGGAATG	1320
GACCACGAGA TTGGTTCCAT CGAAACCGGC AAGCGCGCGG ACCTTATCCT GCTTGACCTG	1380
CGTCACCCTC AGACGACTCC TCACCATCAT TTGGCGGCCA CGATCGTGTT TCAGGCTTAC	1440
GGCAATGAGG TGGACACTGT CCTGATTGAC GGAAACGTTG TGATGGAGAA CCGCCGCTTG	1500
AGCTTTCTTC CCCCTGAACG TGAGTTGGCG TTCCTTGAGG AAGCGCAGAG CCGCGCCACA	1560
GCTATTTTGC AGCGGGCGAA CATGGTGGCT AACCAGCTT GGCGCAGCCT CTAGGAAATG	1620
ACGCCGTTGC TGCATCCGCC GCCCCTTGAG GAAATCGCTG CCATCTTGGC GCGGCTCGGA	1680
TTGGGGGGCG GACATGACCT TGATGGATAC AGAATTGCCA TGAATGCGGC ACTTCCGTCC	1740
TTGCTCGTG TGAATCGTT GGTAGGTGAG GGTGACTGC GGGCGCCAGC TTCCGAAGA	1800
AGTGAAAG	1808

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1846 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GAGCGCCGCC ACAGCAGCCT TGATCATGAA GGCGAGCATG GTGACCTTGA CGCCGCTCTT	60
TTGCTTCTCT TTGTTGAACT GCACGCGAAA GGCTTCCAGG TCGGTGATGT CCGCGTCGTC	120
GTGGTTGGTG ACGTGC GGGA TGACCACCCA GTTGCGGTGC AGGTTTTTCG ATGGCGTAAT	180
ATCTGCGTTG CGACGTGTAA CACACTATTG GAGACATATC ATGCAAACGC TCAGCATCCA	240
GCACGGTACC CTCGTCACGA TGGATCAGTA CCGCAGAGTC CTTGGGGATA GCTGGGTTCA	300
CGTGCAAGAT GGACGGATCG TCGCGCTCGG AGTGACAGCC GAGTCGGTGC CTCCGCCAGC	360

GGATCGGGTG ATCGATGCAC GCGGCAAGGT CGTGTTACCC GGTTCATCA ATGCCCACAC	420
CCATGTGAAC CAGATCCTCC TCGCGGAGG GCCCTCGCAC GGGCGTCAAT TCTATGACTG	480
GCTGTTCAAC GTTGTGTATC CGGGACAAAA GGCGATGAGA CCGGAGGACG TAGCGGTGGC	540
GGTGAGGTTG TATTGTGCGG AAGCTGTGCG CAGCGGGATT ACGACGATCA ACGAAAACGC	600
CGATTCGGCC ATCTACCCAG GCAACATCGA GGCCGCGATG GCGGTCTATG GTGAGGTGGG	660
TGTGAGGGTC GTCTACGCCC GCATGTTCTT TGATCGGATG GACGGGCGCA TTCAAGGGTA	720
TGTGGACGCC TTGAAGGCTC GCTCTCCCCA AGTCGAACTG TGCTCGATCA TGGAGGAAAC	780
GGCTGTGGCC AAAGATCGGA TCACAGCCCT GTCAGATCAG TATCATGGCA CGGCAGGAGG	840
TCGTATATCA GTTTGGCCCG CTCCTGCCAC TACCACGGCG GTGACAGTTG AAGGAATGCG	900
ATGGGCACAA GCCTTCGCCC GTGATCGGGC GGTAATGTGG ACGCTTCACA TGGCGGAGAG	960
CGATCATGAT GAGCGGATTC ATGGGATGAG TCCCGCCGAT TACATGGAGT GTTACGGACT	1020
CTTGATGAG CGTCTGCAGG TCGCGCATTG CGTGACTTT GACCGGAAGG ATGTTCCGGCT	1080
GCTGCACCGC CACAATGTGA AGGTCGCGTC GCAGGTTGTG AGCAATGCCT ACCTCGGCTC	1140
AGGGGTGGCC CCCGTGCCAG AGATGGTGGA GCGCGGCATG GCCGTGGGCA TTGGAACAGA	1200
TAACGGGAAT AGTAATGACT CCGTAAACAT GATCGGAGAC ATGAAGTTTA TGGCCCATAT	1260
TCACCGCGCG GTGCATCGGG ATGCGGACGT GCTGACCCCA GAGAAGATTC TTGAAATGGC	1320
GACGATCGAT GGGGCGCGTT CGTTGGGGAT GGACCACGAG ATTGGTTCCA TCGAAACCGG	1380
CAAGCGCGCG GACCTTATCC TGCTTGACCT GCGTCACCCT CAGACGACTC CTCACCATCA	1440
TTTGGCGGCC ACGATCGTGT TTCAGGCTTA CGGCAATGAG GTGGACACTG TCCTGATTGA	1500
CGGAAACGTT GTGATGGAGA ACCGCCGCTT GAGCTTTCTT CCCCCTGAAC GTGAGTTGGC	1560
GTTCCTTGAG GAAGCGCAGA GCCGCGCCAC AGCTATTTTG CAGCGGGCGA ACATGGTGGC	1620
TAACCCAGCT TGGCGCAGCC TCTAGGAAAT GACGCCGTTG CTGCATCCGC CGCCCCTTGA	1680
GGAAATCGCT GCCATCTTGG CGCGGCTCGG ATTGGGGGGC GGACATGACC TTGATGGATA	1740
CAGAATTGCC ATGAATGCGG CACTTCCGTC CTTGCTCGT GTGGAATCGT TGGTAGGTGA	1800
GGGTCGACTG CGGGCGCCAG CTTCCCGAAG AAGTGAAAGG CCCGAG	1846

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 601 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

42

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Ala	Ser	Met	Val	Thr	Leu	Thr	Pro	Leu	Phe	Ser	Phe	Ser	Leu	Leu	Asn	1	5	10	15
Cys	Thr	Arg	Lys	Ala	Ser	Arg	Ser	Val	Met	Ser	Ala	Ser	Ser	Trp	Leu	20	25	30	
Val	Thr	Cys	Gly	Met	Thr	Thr	Gln	Leu	Arg	Cys	Arg	Phe	Phe	Asp	Gly	35	40	45	
Ile	Ile	Ser	Ala	Leu	Arg	Arg	Val	Thr	His	Tyr	Trp	Arg	His	Ile	Met	50	55	60	
Gln	Thr	Leu	Ser	Ile	Gln	His	Gly	Thr	Leu	Val	Thr	Met	Asp	Gln	Tyr	65	70	75	80
Arg	Arg	Val	Leu	Gly	Asp	Ser	Trp	Val	His	Val	Gln	Asp	Gly	Arg	Ile	85	90	95	
Val	Ala	Leu	Gly	Val	His	Ala	Glu	Ser	Val	Pro	Pro	Pro	Ala	Asp	Arg	100	105	110	
Val	Ile	Asp	Ala	Arg	Gly	Lys	Val	Val	Leu	Pro	Gly	Phe	Ile	Asn	Ala	115	120	125	
His	Thr	His	Val	Asn	Gln	Ile	Leu	Leu	Arg	Gly	Gly	Pro	Ser	His	Gly	130	135	140	
Arg	Gln	Phe	Tyr	Asp	Trp	Leu	Phe	Asn	Val	Val	Tyr	Pro	Gly	Gln	Lys	145	150	155	160
Ala	Met	Arg	Pro	Glu	Asp	Val	Ala	Val	Ala	Val	Arg	Leu	Tyr	Cys	Ala	165	170	175	
Glu	Ala	Val	Arg	Ser	Gly	Ile	Thr	Thr	Ile	Asn	Glu	Asn	Ala	Asp	Ser	180	185	190	
Ala	Ile	Tyr	Pro	Gly	Asn	Ile	Glu	Ala	Ala	Met	Ala	Val	Tyr	Gly	Glu	195	200	205	
Val	Gly	Val	Arg	Val	Val	Tyr	Ala	Arg	Met	Phe	Phe	Asp	Arg	Met	Asp	210	215	220	
Gly	Arg	Ile	Gln	Gly	Tyr	Val	Asp	Ala	Leu	Lys	Ala	Arg	Ser	Pro	Gln	225	230	235	240
Val	Glu	Leu	Cys	Ser	Ile	Met	Glu	Gly	Thr	Ala	Val	Ala	Lys	Asp	Arg	245	250	255	

43

Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly Gly Arg Ile
 260 265 270
 Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr Val Glu Gly
 275 280 285
 Met Arg Trp Ala Gln Ala Phe Ala Arg Asp Arg Ala Val Met Trp Thr
 290 295 300
 Leu His Met Ala Glu Ser Asp His Asp Glu Arg Ile His Gly Met Ser
 305 310 315 320
 Pro Ala Glu Tyr Met Glu Cys Tyr Gly Leu Leu Asp Glu Arg Leu Gln
 325 330 335
 Val Ala His Cys Val Tyr Phe Asp Arg Lys Asp Val Arg Leu Leu His
 340 345 350
 Arg His Asn Val Lys Val Ala Ser Gln Val Val Ser Asn Ala Tyr Leu
 355 360 365
 Gly Ser Gly Val Ala Pro Val Pro Glu Met Val Glu Arg Gly Met Ala
 370 375 380
 Val Gly Ile Gly Thr Asp Asn Gly Asn Ser Asn Asp Ser Val Asn Met
 385 390 395 400
 Ile Gly Asp Met Lys Phe Met Ala His Ile His Arg Ala Val His Arg
 405 410 415
 Asp Ala Asp Val Leu Thr Pro Glu Lys Ile Leu Glu Met Ala Thr Ile
 420 425 430
 Asp Gly Ala Arg Ser Leu Gly Met Asp His Glu Ile Gly Ser Ile Glu
 435 440 445
 Thr Gly Lys Arg Ala Asp Leu Ile Leu Leu Asp Leu Arg His Pro Gln
 450 455 460
 Thr Thr Pro His His His Leu Ala Ala Thr Ile Val Phe Gln Ala Tyr
 465 470 475 480
 Gly Asn Glu Val Asp Thr Val Leu Ile Asp Gly Asn Val Val Met Glu
 485 490 495
 Asn Arg Arg Leu Ser Phe Leu Pro Pro Glu Arg Glu Leu Ala Phe Leu
 500 505 510
 Glu Glu Ala Gln Ser Arg Ala Thr Ala Ile Leu Gln Arg Ala Asn Met
 515 520 525
 Val Ala Asn Pro Ala Trp Arg Ser Leu Glu Met Thr Pro Leu Leu His
 530 535 540
 Pro Pro Pro Leu Glu Glu Ile Ala Ala Ile Leu Ala Arg Leu Gly Leu
 545 550 555 560

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 614 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ser	Ala	Ala	Thr	Ala	Ala	Leu	Ile	Met	Lys	Ala	Ser	Met	Val	Thr	Leu	1	5	10	15
Thr	Pro	Leu	Phe	Ser	Phe	Ser	Leu	Leu	Asn	Cys	Thr	Arg	Lys	Ala	Ser	20	25	30	
Arg	Ser	Val	Met	Ser	Ala	Ser	Ser	Trp	Leu	Val	Thr	Cys	Gly	Met	Thr	35	40	45	
Thr	Gln	Leu	Arg	Cys	Arg	Phe	Phe	Asp	Gly	Val	Ile	Ser	Ala	Leu	Arg	50	55	60	
Arg	Val	Thr	His	Tyr	Trp	Arg	His	Ile	Met	Gln	Thr	Leu	Ser	Ile	Gln	65	70	75	80
His	Gly	Thr	Leu	Val	Thr	Met	Asp	Gln	Tyr	Arg	Arg	Val	Leu	Gly	Asp	85	90	95	
Ser	Trp	Val	His	Val	Gln	Asp	Gly	Arg	Ile	Val	Ala	Leu	Gly	Val	His	100	105	110	
Ala	Glu	Ser	Val	Pro	Pro	Pro	Ala	Asp	Arg	Val	Ile	Asp	Ala	Arg	Gly	115	120	125	
Lys	Val	Val	Leu	Pro	Gly	Phe	Ile	Asn	Ala	His	Thr	His	Val	Asn	Gln	130	135	140	
Ile	Leu	Leu	Arg	Gly	Gly	Pro	Ser	His	Gly	Arg	Gln	Phe	Tyr	Asp	Trp	145	150	155	160
Leu	Phe	Asn	Val	Val	Tyr	Pro	Gly	Gln	Lys	Ala	Met	Arg	Pro	Glu	Asp	165	170	175	

45

Val Ala Val Ala Val Arg Leu Tyr Cys Ala Glu Ala Val Arg Ser Gly
 180 185 190
 Ile Thr Thr Ile Asn Glu Asn Ala Asp Ser Ala Ile Tyr Pro Gly Asn
 195 200 205
 Ile Glu Ala Ala Met Ala Val Tyr Gly Glu Val Gly Val Arg Val Val
 210 215 220
 Tyr Ala Arg Met Phe Phe Asp Arg Met Asp Gly Arg Ile Gln Gly Tyr
 225 230 235 240
 Val Asp Ala Leu Lys Ala Arg Ser Pro Gln Val Glu Leu Cys Ser Ile
 245 250 255
 Met Glu Glu Thr Ala Val Ala Lys Asp Arg Ile Thr Ala Leu Ser Asp
 260 265 270
 Gln Tyr His Gly Thr Ala Gly Gly Arg Ile Ser Val Trp Pro Ala Pro
 275 280 285
 Ala Thr Thr Thr Ala Val Thr Val Glu Gly Met Arg Trp Ala Gln Ala
 290 295 300
 Phe Ala Arg Asp Arg Ala Val Met Trp Thr Leu His Met Ala Glu Ser
 305 310 315 320
 Asp His Asp Glu Arg Ile His Gly Met Ser Pro Ala Asp Tyr Met Glu
 325 330 335
 Cys Tyr Gly Leu Leu Asp Glu Arg Leu Gln Val Ala His Cys Val Tyr
 340 345 350
 Phe Asp Arg Lys Asp Val Arg Leu Leu His Arg His Asn Val Lys Val
 355 360 365
 Ala Ser Gln Val Val Ser Asn Ala Tyr Leu Gly Ser Gly Val Ala Pro
 370 375 380
 Val Pro Glu Met Val Glu Arg Gly Met Ala Val Gly Ile Gly Thr Asp
 385 390 395 400
 Asn Gly Asn Ser Asn Asp Ser Val Asn Met Ile Gly Asp Met Lys Phe
 405 410 415
 Met Ala His Ile His Arg Ala Val His Arg Asp Ala Asp Val Leu Thr
 420 425 430
 Pro Glu Lys Ile Leu Glu Met Ala Thr Ile Asp Gly Ala Arg Ser Leu
 435 440 445
 Gly Met Asp His Glu Ile Gly Ser Ile Glu Thr Gly Lys Arg Ala Asp
 450 455 460

46

Leu Ile Leu Leu Asp Leu Arg His Pro Gln Thr Thr Pro His His His
 465 470 475 480
 Leu Ala Ala Thr Ile Val Phe Gln Ala Tyr Gly Asn Glu Val Asp Thr
 485 490 495
 Val Leu Ile Asp Gly Asn Val Val Met Glu Asn Arg Arg Leu Ser Phe
 500 505 510
 Leu Pro Pro Glu Arg Glu Leu Ala Phe Leu Glu Glu Ala Gln Ser Arg
 515 520 525
 Ala Thr Ala Ile Leu Gln Arg Ala Asn Met Val Ala Asn Pro Ala Trp
 530 535 540
 Arg Ser Leu Glu Met Thr Pro Leu Leu His Pro Pro Pro Leu Glu Glu
 545 550 555 560
 Ile Ala Ala Ile Leu Ala Arg Leu Gly Leu Gly Gly Gly His Asp Leu
 565 570 575
 Asp Gly Tyr Arg Ile Ala Met Asn Ala Ala Leu Pro Ser Phe Ala Arg
 580 585 590
 Val Glu Ser Leu Val Gly Glu Gly Arg Leu Arg Ala Pro Ala Ser Arg
 595 600 605
 Arg Ser Glu Arg Pro Glu
 610

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 545 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGGTATCGGG GAATTCTTGA GCGCGGCCAC AGCAGCCNTG ATCATGAAGG CGAGCATGGT 60
 GACCTNGACG CCGTNTTTTNN GTTNTTTTTTT GTTGAAGTGC ACGCGAAAGG TTCCAGGTGC 120
 GTGATGTCCG CGTCGTCGTG GTTGGTGACG TCGGGGATGA CCACCCAGNT GCGGTGCAGG 180
 TTTTTCGATG GCATAATATC TCGGTTGCGA CGTGTAACAC ACTANTGGAG ACATATCATG 240
 CAAACGCTCA GCATCCAGCA CGGTACCCTC GTCACGATGG ATCAGTACCG CAGAGTCCTT 300
 GGGGATAGCT GGGTTCACGT GCAGGATGGA CGGATCGTCG CGCTCGGAGT GCACGCCGAG 360

47

TCGGTGCCTC CGCCAGCGGA TCGGGTGATC GATGCACGCG GCAAGGTCGT GTTACCCGGT 420
 TTCATCAATG CCCACACCCA TGTGAACCAG ATCCTCCTGC GCGGAGGGCC CTCGCACGGG 480
 CGTCAATTCT ATGACTGGCT GTTCAACGTT GTGTATCCGG GACAAAAGGC GATGAGACCG 540
 GAGGA 545

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 499 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CCTGCGCGGA GGGCCTCCGC ACGGGCGTCA ATTCTATGAC TGGCTGTTCA AC GTTGTGTA 60
 TCCGGGACAA AAGGCGATGA GACCGGAGGA CGTAGCGGTG GCGGTGAGGT TGTATTGTGC 120
 GGAAGCTGTG CGCAGCGGGA TTACGACGAT CAACGAAAAC GCCGATTCGG CCATCTACCC 180
 AGGCAACATC GAGGCCGCGA TGGCGGTCTA TGGTGAGGTG GGTGTGAGGG TCGTCTACGC 240
 CCGCATGTTC TTTGATCGGA TGGACGGGCG CATTCAAGGG TATGTGGACG CCTTGAAGGC 300
 TCGCTCTCCC CAAGTCGAAC TGTGCTCGAT CATGGAGGAA ACGGCTGTGG CCAAAGATCG 360
 GATCACAGCC CTGTCAGATC AGTATCATGG CACGGCAGGA GGTCCTATAT CAGTTTGGCC 420
 CGCTCCTGCC ACTACCACGG CGGTGACATT TAAANGAATC CATGGGCCAA CCTCCCCCGT 480
 GATCCGGCGG TAATGTGAC 499

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 360 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TNGCAGGTTG TGAGCATGCT ACTTCGGTTC AGGNGTGGCC CCCGTGCCAG AGATGGTGGA	60
GCGCGGCATG GCCGTGGGCA TTGGAACAGA TAACGGGAAT AGTAATGACT CCGTAAACAT	120
GATCGGAGAC ATGAAGTTTA TGGCCCATAT TCACCGCGCG GTGCATCGGG ATGCGGACGT	180
GCTGACCCCA GAGAAGATTN TTGAAATGGC GACGATCGAT GGGGCGCGTT TCGTTGGGGA	240
TGGACCACGA GATTGGTTCC ATCGAAACCG GCAAGCGCGC GGACCTTATC CTGCTTGACC	300
TGCGTCACCC TCAGACGACT CCTCACCATC ATTTGGCGGC CACGATCGTG TTTCAGGCTT	360

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 443 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CGGCCACGAT CGTGTTCAG GCTTACGGCA ATGAGGTGGA CACTGTCCTG ATTGACGGAA	60
ACGTTGTGAT GGAGAACCGC CGCTTGAGCT TTCTTCCCCC TGAACGTGAG TTGGCGTTCC	120
TTGAGGAAGC GCAGAGCCGC GCCACAGCTA TTTTGCATCG GGCGAAACAT GGTGGCTAAC	180
CCAGCTTGGC GCAGCCTCTA GGAAATGACG CCGTTGCTGC ATCCGCCGCC CCTTGAGGAA	240
ATCGCTGCCA TCTTGGCGCG GCTCGGATTG GGGGGCGGAC ATGACCTTGA TGGATACAGA	300
ATTGCCATGA ATGCGGCACT TCCGTCCTTC GCTCGTGTGG AATCGTTGGT AGGTGAGGGT	360
CGACTGCGGG CGCCAGCTTC CCGAAGAGGT GAAAGCCCGA GGATCCTCTA GAGTCCGATT	420
TTTCCGATGT CATCACCGGC GCG	443

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 505 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

```

CCTGCGCGGA GGCCTCCGCA CGGGCGTCAA TTCTATGACT GGCTGTTCAA CGTTGTGTAT      60
CCGGGACAAA AGGCGATGAG ACCGGAGGAC GTANCGGTGG CGGTGAGGTT GTATTGTGCG      120
GAAGCTGTGC GCAGCGGGAT TACGACGATC AACGAAAACG CCGATTCGGC CATCTACCCA      180
GGCAACATCG AGGCCGCGAT GGCGGTCTAT GGTGAGGTGG GTGTGAGGGT CGTCTACGCC      240
CGCATGTTCT TTGATCGGAT GGACGGGCGC ATTCAAGGGT ATGTGGACGC CTTGAAGGCT      300
CGCTCTCCCC AAGTCGAACT GTGCTCGATC ATGGAGGAAA CGGCTGTGGC CAAAGATCGG      360
ATCACANCCC TGTCAGATCA NTATCATGGC ACGGCANGAG GTCCTATATC ANTTTGGCCC      420
GTCCTGCCA CTACCACNGC GGTGACATTT NAANGAATTC CATNGGCACA ACCTTCCCCC      480
GTGATCNGGC GGTAATGTNG ACCCA                                           505

```

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 144 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

Pro His Gly Arg Gln Phe Tyr Asp Trp Leu Phe Asn Val Leu Tyr Pro
1           5           10           15
Gly Gln Lys Ala Met Arg Pro Glu Asp Val Ala Val Ala Val Arg Leu
20           25           30
Tyr Cys Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn
35           40           45
Ala Asp Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val
50           55           60
Tyr Gly Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp
65           70           75           80
Arg Met Asp Gly Arg Ile Gln Gly Tyr Val Asp Ala Leu Lys Ala Arg
85           90           95
Ser Pro Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val Ala
100          105          110

```

50

Lys Asp Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly
 115 120 125

Gly Arg Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr
 130 135 140

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 144 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Ser His Gly Arg Gln Phe Tyr Asp Trp Leu Phe Asn Val Leu Tyr Pro
 1 5 10 15

Gly Gln Lys Ala Met Arg Pro Glu Asp Val Ala Val Ala Val Arg Leu
 20 25 30

Tyr Cys Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn
 35 40 45

Ala Asp Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val
 50 55 60

Tyr Gly Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp
 65 70 75 80

Arg Met Asp Gly Arg Ile Gln Gly Tyr Val Asp Thr Leu Lys Ala Arg
 85 90 95

Ser Pro Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val Ala
 100 105 110

Lys Asp Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly
 115 120 125

Gly Arg Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr
 130 135 140

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 144 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single

51

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```

Pro His Gly Arg Gln Phe Tyr Asp Trp Leu Phe Asn Val Val Tyr Pro
1           5           10           15
Gly Gln Lys Ala Met Arg Pro Glu Asp Val Ala Val Ala Val Arg Leu
20           25           30
Tyr Cys Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn
35           40           45
Ala Asp Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val
50           55           60
Tyr Gly Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp
65           70           75           80
Arg Met Asp Gly Arg Ile Gln Gly Tyr Val Asp Ala Leu Lys Ala Arg
85           90           95
Ser Pro Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val Ala
100          105          110
Lys Asp Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly
115          120          125
Gly Arg Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr
130          135          140

```

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 145 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

```

Ser His Gly Arg Gln Phe Tyr Asp Trp Leu Phe Asn Val Leu Tyr Pro
1           5           10           15

```

52

Gly Gln Lys Ala Met Arg Pro Glu Asp Val Ala Val Ala Val Arg Leu
 20 25 30
 Tyr Cys Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn
 35 40 45
 Asn Ala Asp Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala
 50 55 60
 Val Tyr Gly Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe
 65 70 75 80
 Asp Arg Met Asp Gly Arg Ile Gln Gly Tyr Val Asp Thr Leu Lys Ala
 85 90 95
 Arg Ser Pro Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val
 100 105 110
 Ala Lys Asp Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala
 115 120 125
 Gly Gly Arg Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val
 130 135 140
 Thr
 145

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 144 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Ser His Gly Arg Gln Phe Tyr Asp Trp Leu Phe Asn Val Val Tyr Pro
 1 5 10 15
 Gly Gln Lys Ala Met Arg Pro Glu Asp Val Ala Val Ala Val Arg Leu
 20 25 30
 Tyr Cys Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn
 35 40 45
 Ala Asp Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val
 50 55 60
 Tyr Gly Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp
 65 70 75 80

Arg	Met	Asp	Gly	Arg	Ile	Gln	Gly	Tyr	Val	Asp	Ala	Leu	Lys	Ala	Arg
				85					90					95	
Ser	Pro	Gln	Val	Glu	Leu	Cys	Ser	Ile	Met	Glu	Glu	Thr	Ala	Val	Ala
			100					105					110		
Lys	Asp	Arg	Ile	Thr	Ala	Leu	Ser	Asp	Gln	Tyr	His	Gly	Thr	Ala	Gly
		115					120					125			
Gly	Arg	Ile	Ser	Val	Trp	Pro	Ala	Pro	Ala	Thr	Thr	Thr	Ala	Val	Thr
	130					135					140				

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1633 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CGCGAAAGGC	TTCCAGGTCG	GTGATGTCCG	CGTCGTCGTG	GTTGGTGACG	TGCGGGATGA	60
CCACCCAGTC	GCGGTGCAGG	TTTTTTCGATG	GCATAATATC	TGCGTTGCGA	CGTGTAACAC	120
ACTATTGGAG	ACATATCATG	CAAACGCTCA	GCATCCAGCA	CGGTACCCTC	GTCACGATGG	180
ATCAATACCG	CAGAGTCCTT	GGGGATAGCT	GGGTTCACGT	GCAGGATGGA	CGGATCGTCG	240
CGCTCGGAGT	GCACGCCAAG	TCGGTGCCTC	CGCCAGCGGA	TCGGGTGATC	GATGCACGCG	300
GCAAGGTCGT	GTTACCCGGT	TTCATCAATG	CCCACACCCA	TGTGAACCAG	ATCCTCCTGC	360
GCGGAGGGCC	CTCGCACGGG	CGTCAATTCT	ATGACTGGCT	GTTCAACGTT	GTGTATCCGG	420
GACAAAAGGC	GATGAGACCG	GAGGACGTAG	CGGTGGCGGT	GAGGTTGTAT	TGTGCGGAAG	480
CTGTGCGCAG	CGGGATTACG	ACGATCAACG	AAAACGCCGA	TTCGGCCATC	TACCCAGGCA	540
ACATCGAGGC	CGCGATGGCG	GTCTATGGTG	AGGTGGGTGT	GAGGGTCGTC	TACGCCC GCA	600
TGTTCTTTGA	TCGGATGGAC	GGGCGCATTG	AAGGGTATGT	GGACGCCTTG	AAGGCTCGCT	660
CTCCCCAAGT	CGAACTGTGC	TCGATCATGG	AGGAAACGGC	TGTGGCCAAA	GATCGGATCA	720
CAGCCCTGTC	AGATCAGTAT	CATGGCACGG	CAGGAGGTCG	TATATCAGTT	TGGCCCGCTC	780
CTGCCACTAC	CACGGCGGTG	ACAGTTGAAG	GAATGCGATG	GGCACAAGCC	TTCGCCCCGTG	840

ATCGGGCGGT AATGTGGACG CTTCACATGG CGGAGAGCGA TCATGATGGG CGGATTCATG	900
GGATGAGTCC CGCCGAGTAC ATGGAGTGTT ACGGACTCTT GGATGAGCGT CTGCAGGTCC	960
CGCATTGCGT GTACTTTGAC CGGAAGGATG TTCGGCTGCT GCACCGCCAC AATGTGAAGG	1020
TCGCGTCGCA GGTGTGAGC AATGCCTACC TCGGCTCAGG GGTGGCCCCC GTGCCAGAGA	1080
TGGTGGAGCG CGGCATGGCC GTGGGCATTG GAACAGATAA CGGGAATAGT AATGACTCCG	1140
TAAACATGAT CGGAGACATG AAGTTTATGG CCCATATTCA CCGCGCGGTG CATCGGGATG	1200
CGGACGTGCT GACCCAGAG AAGATTCTTG AAATGGCGAC GATCGATGGG GCGCGTTCGT	1260
TGGGGATGGA CCACGAGATT GGTTCATCG AAACCGGCAA GCGCGCGGAC CTTATCCTGC	1320
TTGACCTGCG TCACCCTCAG ACGACTCCTC ACCATCATTT GCGGCCACG ATCGTGTTC	1380
AGGCTTACGG CAATGAAGTG GACACTGTCC TGATTGACGG AAACGTTGTG ATGGAGAACC	1440
GCTGCTTGAG CTTTCTTCCC CCTGAACGTG AGTTGGCGTT CCTTGAGGGA GCGCAGAGCC	1500
GCGCCACAGC TATTTTGCAG CGGGCGAACA TGGTGGCTAA CCCAGCTTGG CGCAGCCTCT	1560
AGGAAATGAC GCCGTTGCTG CATCCGCCGC CCCTTGAGGA AATCGCTGCC ATCTTGGCGC	1620
GGCTCGGATT GGG	1633

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1598 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

TCGTGGTTGG TGACGTGCGG GATGACCACC CAGTCGCGGT GCAGGTTTTT CGATGGCATA	60
ATATCTGCGT TGCGACGTGT AACACACTAT TGGAGACATA TCATGCAAAC GCTCAGCATC	120
CAGCACGGTA CCCTCGTCAC GATGGATCAG TACCGCAGAG TCCTTGGGGA TAGCTGGGTT	180
CACGTGCAGG ATGGACGGAT CGTCGCGCTC GGAGTGCACG CCGAGTCGGT GCCTCCGCCA	240
GCGGATCGGG TGATCGATGC ACGCGGCAAG GTCGTGTTAC CCGGTTTCAT CAATGCCAC	300
ACCCATGTGA ACCAGATCCT CCTGCGCGGA GGGCCCTCGC ACGGGCGTCA ATTCTATGAC	360
TGGCTGTTCA ACGTTGTGTA TCCGGGACAA AAGGCGATGA GACCGGAGGA CGTAGCGGTG	420

GCGGTGAGGT TGTATTGTGC GGAAGCTGTG CGCAGCGGGA TTACGACGAT CAACGAAAAC	480
GCCGATTCGG CCATCTACCC AGGCAACATC GAGGCCGCGA TGGCGGTCTA TGGTGAGGTG	540
GGTGTGAGGG TCGTCTACGC CCGCATGTTC TTTGATCGGA TGGACGGGCG CATTCAAGGG	600
TATGTGGACG CCTTGAAGGC TCGCTCTCCC CAAGTCGAAC TGTGCTCGAT CATGGAGGAA	660
ACGGCTGTGG CCAAAGATCG GATCACAGCC CTGTCAGATC AGTATCATGG CACGGCAGGA	720
GGTCGTATAT CAGTTTGGCC CGCTCCTGCC ACTACCACGG CGGTGACAGT TGAAGGAATG	780
CGATGGGCAC AAGCCTTCGC CCGTGATCGG GCGGTAATGT GGACGCTTCA CATGGCGGAG	840
AGCGATCATG ATGAGCGGAT TCATGGGATG AGTCCC GCCG AGTACATGGA GTGTCACGGA	900
CTCTTGATG AGCGTCTGCA GGTGCGCAT TGC GTACT TTGACCGGAA GGATGTTCCG	960
CTGCTGCACC GCCACAATGT GAAGGTCGCG TCGCAGGTTG TGAGCAATGC CTACCTCGGC	1020
TCAGGGGTGG CCCCCGTGCC AGAGATGGTG GAGCGCGGCA TGGCCATGGG CATTGGAACA	1080
GATAACGGGA ATAGTAATGA CTCCGTAAAC ATGATCGGAG ACATGAAGTT TATGGCCCAT	1140
ATTCACCGCG CGGTGCATCG GGATGCGGAC GTGCTGACCC CAGAGAAGAT TCTTGAAATG	1200
GCGACGATCG ATGGGGCGCG TTCGTTGGGA ATGGACCACG AGATTGGTTC CATCGAAACC	1260
GGCAAGCGCG CGGACCTTAT CCTGCTTGAC CTGCGTCACC CTCAGACGAC TCCTCACCAT	1320
CATTGCGCG CCACGATCGT GTTTCAGGCT TACGGCAATG AGGTGGACAC TGTCTGATT	1380
GACGGAACG TTGTGATGGA GAACCGCCGC TTGAGCTTTC TTCCCCCTGA ACGTGAGTTG	1440
GCGTTCCTTG AGGAAGCGCA GAGCCGCGCC ACAGCTATTT TGCAGCGGGC GAACATGGTG	1500
GCTAACCAG CTTGGCGCAG CCTCTAGGAA ATGACGCCGT TGCTGCATCC GCCGCCCTT	1560
GAGGAAATCG CTGCCATCTT GGCGCGGCTC GGATTGGG	1598

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1586 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

ACGTGCGGGA TGACCACCA GTTGC GTGC AGGTTTTTCG ATGGCGTAAT ATCTGCGTTG 60

CGACGTGTAA CACACTATTG GAGACATATC ATGCAAACGC TCAGCATCCA GCACGGTACC	120
CTCGTCACGA TGGATCAGTA CCGCAGAGTC CTTGGGGATA GCTGGGTTCG CGTGCAGGAT	180
GGACGGATCG TCGCGCTCGG AGTGCACGCC GAGTCGGTGC CTCCGCCAGC GGATCGGGTG	240
ATCGATGCAC GCGGCAAGGT CGTGTTACCC GGTTCATCA ATGCCACAC CCATGTGAAC	300
CAGATCCTCC TGC GCGGAGG GCCCTCGCAC GGGCGTCAAT TCTATGACTG GCTGTTCAAC	360
GTTGTGTATC CGGGACAAAA GGCGATGAGA CCTGAGGACG TAGCGGTGGC GGTGAGGTTG	420
TATTGTGCGG AAGCTGTGCG CAGCGGGATT ACGACGATCA ACGAAAACGC CGATTCGGCC	480
ATCTACCCAG GCAACATCGA GGCCGCGATG GCGGTCTATG GTGAGGTGGG TGTGAGGGTC	540
GTCTACGCCC GCATGTTCTT TGATCGGATG GACGGGCGCA TTCAAGGGTA TGTGGACGCC	600
TTGAAGGCTC GCTCTCCCCA AGTCGAACTG TGCTCGATCA TGGAGGAAAC GGCTGTGGCC	660
AAAGATCGGA TCACAGCCCT GTCAGATCAG TATCATGGCA CGGCAGGAGG TCGTATATCA	720
GTTTGGCCCC CTCCTGCCAC TACCACGGCG GTGACAGTTG AAGGAATGCG ATGGGCACAA	780
GCCTTCGCCC GTGATCGGGC GGTAATGTGG ACGCTTCACA TGGCGGAGAG CGATCATGAT	840
GAGCGGATTC ATGGGATGAG TCCCGCCGAG TACATGGAGT GTTACGGACT CTTGGATGAG	900
CGTCTGCAGG TCGCGCATTG CGTGTACTTT GACCGGAAGG ATGTTCCGGCT GCTGCACCGC	960
CACAATGTGA AGGTGCGGTC GCAGGTTGTG AGCAATGCCT ACCTCGGCTC AGGGGTGGCC	1020
CCCGTGCCAG AGATGGTGGG GCGCGGCATG GCCGTGGGCA TTGGAACAGA TAACGGGAAT	1080
AGTAATGACT CCGTAAACAT GATCGGAGAC ATGAAGTTTA TGGCCCATAT TCACCGCGCG	1140
GTGCATCGGG ATGCGGACGT GCTGACCCCA GAGAAGATTC TTGAAATGGC GACAATCGAT	1200
GGGGCGCGTT CGTTGGGAAT GGACCACGAG ATTGGTTCCA TCGAAACCGG CAAGCGCGCG	1260
GACCTTATCC TGCTTGACCT GCGTCACCCT CAGACGACTC CTCACCATCA TTTGGCGGCC	1320
ACGATCGTGT TTCAGGCTTA CGGCAATGAG GTGGACACTG TCCTGATTGA CGGAAACGTT	1380
GTGATGGAGA ACCGCCGCTT GAGCTTTCTT CCCCCTGAAC GTGAGTTGGC GTTCCTTGAG	1440
GAAGCGCAGA GCCGCGCCAC AGCTATTTTG CAGCGGGCGA ACATGGTGGC TAACCCAGCT	1500
TGGCGCAGCC TCTAGGAAAT GACGCCGTTG CTGCATCCGC TGCCCCCTGA GGAAATCGCT	1560
GCCATCTTGG CGCGGCTCGG ATTGGG	1586

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1597 base pairs

- (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

CGTGGTTGGT GACGTGGGGG ATGACCACCC AGTCGCGGTG CAGGTTTTTC GATGGCATAA	60
TATCTGCGTT GCGACGTGTA ACACACTATT GGAGACATAT CATGCAAACG CTCAGCATCC	120
AGCACGGTAC CCTCGTCACG ATGGATCAGT ACCGCAGAGT CCTTGGGGAT AGCTGGGTTT	180
ACGTGCAGGA TGGACGGATC GTCGCGCTCG GAGTGCACGC CGAGTCGGTG CCTCCGCCAG	240
CGGATCAGGT GATCGATGCA CGCGGCAAGG TCGTGTTACC CGGTTTCATC AATGCCCACA	300
CCCATGTGAA CCAGATCCTC CTGCGCGGAG GGCCCTCGCA CGGGCGTCAA TTCCATGACT	360
GGCTGTTCAA CGTTGTGTAT CCGGGACAAA AGGCGATGAG ACCGGAGGAC GTAGCGGTGG	420
CGGTGAGGTT GTATTGTGCA GAAGCTGTGC GCAGCGGGAT TACGACGATT AACGAAAACG	480
CCGATTCGGC CATCTACCCA GGCAACATCG AGGCCGCGAT GGCGGTCTAT GGTGAGGTGG	540
GTGTGAGGGT CGTCTACGCC CGCATGTTCT TTGATCGGAT GGACGGGCGC ATTCAAGGGT	600
ATGTGGACGC CTTGAAGGCT CGCTCTCCCC AAGTCGAAC TGTCTCGATC ATGGAGGAAA	660
CGGCTGTGGC CAAAGATCGG ATCACAGCCC TGTCAGATCA GTATCATGGC ACGGCAGGAG	720
GTCGTATATC AGTTTGGCCC GTCCTGCCA CTACCACGGC GGTGACAGTT GAAGGAATGC	780
GATGGGCACA AGCCTTCGCC CGTGATCGGG CGGTAATGTG GACGCTTCAC ATGGCGGAGA	840
GCGATCATGA TGGGCGGATT CATGGGATGA GTCCCGCCGA GTACATGGAG TGTACGGAC	900
TCTTGATGA GCGTCTGCAG GTCGCGCATT GCGTGTACTT TGACCGGAAG GATGTTCCGC	960
TGCTGCACCG CCACAATGTG AAGGTCGCGT CGCAGGTTGT GAGCAATGCC TACCTCGGCT	1020
CAGGGGTGGC CCCCCTGCCA GAGATGGTGG AGCGCGGCAT GGCCGTGGGC ATTGGAACAG	1080
ATAACGGGAA TAGTAATGAC TCCGTAAACA TGATCGGAGA CATGAAGTTT ATGGCCCATA	1140
TTCACCGCGC GGTGCATCGG GATGCGGACG TGCTGACCCC AGAGAAGATT CTTGAAATGG	1200
CAACGATCGA TGGGGCGCGT TCGTTGGGAA TGGACCACGA GATTGGTTCC ATCGAAACCG	1260
GCAAGCGCGC GGACCTTATC CTGCTTGACC TGCCTCACCC TCAGACGACT CCTCACCATC	1320
ATTTGGCGGC CACGATCGTG TTTCAGGCTT ACGGCAATGA GGTGGACACT GTCCTGATTG	1380

ACGGAAACGT TGTGATGGAG AACCGCCGCT TGAGCTTTCT TCCCCCTGAA CGTGAGTTGG	1440
CGTTCCTTGA GGAAGCGCAG AGCCGCGCCA CAGCTATTTT GCAGCGGGCG AACATGGTGG	1500
CTAACCCAGC TTGGCGCAGC CTCTAGGAAA TGACGCCGTT GCTGCATCCG CCGCCCCTTG	1560
AGGAAATCGC TGCCATCTTG GCGCGGCTCG GATTGGG	1597

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1674 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GTGACCTTGA CGCCGCTCTT TTCGTTCTCT TTGTTGAACT GCACGCGAAT GGCTTCCAGT	60
TCGATGATGT CCGCGTCGTC GTGGTTGGTG ACGTGCGGGA TGACCACCCA GTCGCGGTGC	120
AGGTTTTTCG ATGGCATAAT ATCTGCGTTG CGACGTGTAA CACACTATTG GAGACATATC	180
ATGCAAACGC TCAGCATCCA GCACGGTACC CTCGTCACGA TGGATCAGTA CCGCAGAGTC	240
CTTGGGGATA GCTGGGTTCG CGTGCAGGAT GGACGGATCG TCGCGCTCGG AGTGCACGCC	300
GAGTCGGTGC CTCCGCCAGC GGATCGGGTG ATTGATGCAC GCGGCAAGGT CGTGTTACCC	360
GGTTTCATCA ATGCCACAC CCATGTGAAC CAGATCCTCC TGCSCGGAGG CCTCGCACGG	420
GCGTCAATTC TATGACTGGC TGTTC AACGT TGTGTATCCG GGACAAAAGG CGATGAGACC	480
GGAGGACGTA GCGGTGGCGG TGAGGTTGTA TTGTGCGGAA GCTGTGCGCA GCGGGATTAC	540
GACGATCAAC GAAAACGCCG ATTCGGCCAT CTACCCAGGC AACATCGAGG CCGCGATGGC	600
GGTCTATGGT GAGGTGGGTG TGAGGGTCGT CTACGCCCGC ATGTTCTTTG ATCGGATGGA	660
CAGGCGCATT CAAGGGTATG TGGACGCCTT GAAGGCTCGC TCTCCCCAAG TCGAACTGTG	720
CTCGATCATG GAGGAAACGG CTGTGGCCAA AGATCGGATC ACAGCCCTGT CAGATCAGTA	780
TCATGGCACG GCAGGAGGTC GTATATCAGT TTGGCCCCTT CCTGCCACTA CCACGGCGGT	840
GACAGTTGAA GGAATGCGAT GGGCACAAGC CTTGCCCCGT GATCGGGCGG TAATGTGGAC	900
GCTTCACATG GCGGAGAGCG ATCATGATGA GCGGATTCAT GGGATGAGTC CCGCCGAGTA	960
CATGGAGTGT TACGGA CTCT TGGATGAGCG TCTGCAGGTC GCGCATTGCG TGTACTTTGA	1020

```

CCGGAAGGAT ATTCGGCTGC TGCACCGCCA CAATGTGAAG GTCGCGTCGC AGGCTGTGAG      1080
CAATGCCTAC CTCGGCTCAG GGGTGGCCCC CGTGCCAGAG ATGGTGGAGC GCGGCATGGC      1140
CGTGGGCATT GGAACAGATA ACGGGAATAG TAATGACTCC GTAAACATGA TCGGAGACAT      1200
GAAGTTTATG GCCCATATTC ACCGCGCGGT GCATCGGGAT GCGGACGTGC TGACCCCAAGA      1260
GAAGATTCTT GAAATGGCGA CGATCGATGG GGCGCGTTTC TTGGGAATGG ACCACGAGAT      1320
TGGTTCATC GAAACCGCA AGCGCGCGGA CCTTATCCTG CTTGACCTGC GTCACCCTCA      1380
GACGACTCCT CACCATCATT TGGCGGCCAC GATCGTGTTC CAGGCTTACG GCAATGAGGT      1440
GGACACTGTC CTGATTGACG GAAACGTTGT GATGGAGAAC CGCCGCTTGA GCTTTCTTCC      1500
CCCTGAACGT GAGTTGGCGT TCCTTGAGGA AGCGCAGAGC CGCGCCACAG CTATTTTGCA      1560
GCGGGCGAAC ATGGTGGCCA ACCCAGCTTG GCGCAGCCTC TAGGAAATGA CGCCGTTGCT      1620
GCATCCGCCG CCCCTTGAGG AAATCGCTGC CATCTTGGCG CAGCTCGGAT TGGG      1674

```

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

```

Met Gln Thr Leu Ser Ile Gln His Gly Thr Leu Val Thr Met Asp Gln
1           5           10           15
Tyr Arg Arg Val Leu Gly Asp Ser Trp Val His Val Gln Asp Gly Arg
20           25           30
Ile Val Ala Leu Gly Val His Ala Lys Ser Val Pro Pro Pro Ala Asp
35           40           45
Arg Val Ile Asp Ala Arg Gly Lys Val Val Leu Pro Gly Phe Ile Asn
50           55           60
Ala His Thr His Val Asn Gln Ile Leu Leu Arg Gly Gly Pro Ser His
65           70           75           80
Gly Arg Gln Phe Tyr Asp Trp Leu Phe Asn Val Val Tyr Pro Gly Gln
85           90           95

```

60

Lys Ala Met Arg Pro Glu Asp Val Ala Val Ala Val Arg Leu Tyr Cys
 100 105 110
 Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn Ala Asp
 115 120 125
 Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val Tyr Gly
 130 135 140
 Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp Arg Met
 145 150 155 160
 Asp Gly Arg Ile Gln Gly Tyr Val Asp Ala Leu Lys Ala Arg Ser Pro
 165 170 175
 Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val Ala Lys Asp
 180 185 190
 Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly Gly Arg
 195 200 205
 Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr Val Glu
 210 215 220
 Gly Met Arg Trp Ala Gln Ala Phe Ala Arg Asp Arg Ala Val Met Trp
 225 230 235 240
 Thr Leu His Met Ala Glu Ser Asp His Asp Gly Arg Ile His Gly Met
 245 250 255
 Ser Pro Ala Glu Tyr Met Glu Cys Tyr Gly Leu Leu Asp Glu Arg Leu
 260 265 270
 Gln Val Ala His Cys Val Tyr Phe Asp Arg Lys Asp Val Arg Leu Leu
 275 280 285
 His Arg His Asn Val Lys Val Ala Ser Gln Val Val Ser Asn Ala Tyr
 290 295 300
 Leu Gly Ser Gly Val Ala Pro Val Pro Glu Met Val Glu Arg Gly Met
 305 310 315 320
 Ala Val Gly Ile Gly Thr Asp Asn Gly Asn Ser Asn Asp Ser Val Asn
 325 330 335
 Met Ile Gly Asp Met Lys Phe Met Ala His Ile His Arg Ala Val His
 340 345 350
 Arg Asp Ala Asp Val Leu Thr Pro Glu Lys Ile Leu Glu Met Ala Thr
 355 360 365
 Ile Asp Gly Ala Arg Ser Leu Gly Met Asp His Glu Ile Gly Ser Ile
 370 375 380
 Glu Thr Gly Lys Arg Ala Asp Leu Ile Leu Leu Asp Leu Arg His Pro
 385 390 395 400

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Met	Gln	Thr	Leu	Ser	Ile	Gln	His	Gly	Thr	Leu	Val	Thr	Met	Asp	Gln	
1				5					10					15		
Tyr	Arg	Arg	Val	Leu	Gly	Asp	Ser	Trp	Val	His	Val	Gln	Asp	Gly	Arg	
			20					25					30			
Ile	Val	Ala	Leu	Gly	Val	His	Ala	Glu	Ser	Val	Pro	Pro	Pro	Ala	Asp	
		35					40					45				
Arg	Val	Ile	Asp	Ala	Arg	Gly	Lys	Val	Val	Leu	Pro	Gly	Phe	Ile	Asn	
	50					55					60					
Ala	His	Thr	His	Val	Asn	Gln	Ile	Leu	Leu	Arg	Gly	Gly	Pro	Ser	His	
65					70					75					80	
Gly	Arg	Gln	Phe	Tyr	Asp	Trp	Leu	Phe	Asn	Val	Val	Tyr	Pro	Gly	Gln	
				85					90					95		
Lys	Ala	Met	Arg	Pro	Glu	Asp	Val	Ala	Val	Ala	Val	Arg	Leu	Tyr	Cys	
			100					105						110		

Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn Ala Asp
 115 120 125
 Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val Tyr Gly
 130 135 140
 Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp Arg Met
 145 150 155 160
 Asp Gly Arg Ile Gln Gly Tyr Val Asp Ala Leu Lys Ala Arg Ser Pro
 165 170 175
 Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val Ala Lys Asp
 180 185 190
 Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly Gly Arg
 195 200 205
 Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr Val Glu
 210 215 220
 Gly Met Arg Trp Ala Gln Ala Phe Ala Arg Asp Arg Ala Val Met Trp
 225 230 235 240
 Thr Leu His Met Ala Glu Ser Asp His Asp Glu Arg Ile His Gly Met
 245 250 255
 Ser Pro Ala Glu Tyr Met Glu Cys His Gly Leu Leu Asp Glu Arg Leu
 260 265 270
 Gln Val Ala His Cys Val Tyr Phe Asp Arg Lys Asp Val Arg Leu Leu
 275 280 285
 His Arg His Asn Val Lys Val Ala Ser Gln Val Val Ser Asn Ala Tyr
 290 295 300
 Leu Gly Ser Gly Val Ala Pro Val Pro Glu Met Val Glu Arg Gly Met
 305 310 315 320
 Ala Met Gly Ile Gly Thr Asp Asn Gly Asn Ser Asn Asp Ser Val Asn
 325 330 335
 Met Ile Gly Asp Met Lys Phe Met Ala His Ile His Arg Ala Val His
 340 345 350
 Arg Asp Ala Asp Val Leu Thr Pro Glu Lys Ile Leu Glu Met Ala Thr
 355 360 365
 Ile Asp Gly Ala Arg Ser Leu Gly Met Asp His Glu Ile Gly Ser Ile
 370 375 380
 Glu Thr Gly Lys Arg Ala Asp Leu Ile Leu Leu Asp Leu Arg His Pro
 385 390 395 400

Gln	Thr	Thr	Pro	His	His	Leu	Ala	Ala	Thr	Ile	Val	Phe	Gln	Ala	
				405					410					415	
Tyr	Gly	Asn	Glu	Val	Asp	Thr	Val	Leu	Ile	Asp	Gly	Asn	Val	Val	Met
			420					425					430		
Glu	Asn	Arg	Arg	Leu	Ser	Phe	Leu	Pro	Pro	Glu	Arg	Glu	Leu	Ala	Phe
		435					440					445			
Leu	Glu	Glu	Ala	Gln	Ser	Arg	Ala	Thr	Ala	Ile	Leu	Gln	Arg	Ala	Asn
	450					455					460				
Met	Val	Ala	Asn	Pro	Ala	Trp	Arg	Ser	Leu	Glu	Met	Thr	Pro	Leu	Leu
465					470					475					480
His	Pro	Pro	Pro	Leu	Glu	Glu	Ile	Ala	Ala	Ile	Leu	Ala	Arg	Leu	Gly
				485					490					495	

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Met	Gln	Thr	Leu	Ser	Ile	Gln	His	Gly	Thr	Leu	Val	Thr	Met	Asp	Gln
1			5						10					15	
Tyr	Arg	Arg	Val	Leu	Gly	Asp	Ser	Trp	Val	His	Val	Gln	Asp	Gly	Arg
			20					25					30		
Ile	Val	Ala	Leu	Gly	Val	His	Ala	Glu	Ser	Val	Pro	Pro	Pro	Ala	Asp
		35					40					45			
Arg	Val	Ile	Asp	Ala	Arg	Gly	Lys	Val	Val	Leu	Pro	Gly	Phe	Ile	Asn
	50					55					60				
Ala	His	Thr	His	Val	Asn	Gln	Ile	Leu	Leu	Arg	Gly	Gly	Pro	Ser	His
65					70					75					80
Gly	Arg	Gln	Phe	Tyr	Asp	Trp	Leu	Phe	Asn	Val	Val	Tyr	Pro	Gly	Gln
			85						90					95	
Lys	Ala	Met	Arg	Pro	Glu	Asp	Val	Ala	Val	Ala	Val	Arg	Leu	Tyr	Cys
			100					105					110		

Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn Ala Asp
 115 120 125
 Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val Tyr Gly
 130 135 140
 Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp Arg Met
 145 150 155 160
 Asp Gly Arg Ile Gln Gly Tyr Val Asp Ala Leu Lys Ala Arg Ser Pro
 165 170 175
 Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val Ala Lys Asp
 180 185 190
 Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly Gly Arg
 195 200 205
 Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr Val Glu
 210 215 220
 Gly Met Arg Trp Ala Gln Ala Phe Ala Arg Asp Arg Ala Val Met Trp
 225 230 235 240
 Thr Leu His Met Ala Glu Ser Asp His Asp Glu Arg Ile His Gly Met
 245 250 255
 Ser Pro Ala Glu Tyr Met Glu Cys Tyr Gly Leu Leu Asp Glu Arg Leu
 260 265 270
 Gln Val Ala His Cys Val Tyr Phe Asp Arg Lys Asp Val Arg Leu Leu
 275 280 285
 His Arg His Asn Val Lys Val Ala Ser Gln Val Val Ser Asn Ala Tyr
 290 295 300
 Leu Gly Ser Gly Val Ala Pro Val Pro Glu Met Val Glu Arg Gly Met
 305 310 315 320
 Ala Val Gly Ile Gly Thr Asp Asn Gly Asn Ser Asn Asp Ser Val Asn
 325 330 335
 Met Ile Gly Asp Met Lys Phe Met Ala His Ile His Arg Ala Val His
 340 345 350
 Arg Asp Ala Asp Val Leu Thr Pro Glu Lys Ile Leu Glu Met Ala Thr
 355 360 365
 Ile Asp Gly Ala Arg Ser Leu Gly Met Asp His Glu Ile Gly Ser Ile
 370 375 380
 Glu Thr Gly Lys Arg Ala Asp Leu Ile Leu Leu Asp Leu Arg His Pro
 385 390 395 400
 Gln Thr Thr Pro His His His Leu Ala Ala Thr Ile Val Phe Gln Ala
 405 410 415

65

Tyr Gly Asn Glu Val Asp Thr Val Leu Ile Asp Gly Asn Val Val Met
 420 425 430
 Glu Asn Arg Arg Leu Ser Phe Leu Pro Pro Glu Arg Glu Leu Ala Phe
 435 440 445
 Leu Glu Glu Ala Gln Ser Arg Ala Thr Ala Ile Leu Gln Arg Ala Asn
 450 455 460
 Met Val Ala Asn Pro Ala Trp Arg Ser Leu Glu Met Thr Pro Leu Leu
 465 470 475 480
 His Pro Leu Pro Leu Glu Glu Ile Ala Ala Ile Leu Ala Arg Leu Gly
 485 490 495

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Met Gln Thr Leu Ser Ile Gln His Gly Thr Leu Val Thr Met Asp Gln
 1 5 10 15
 Tyr Arg Arg Val Leu Gly Asp Ser Trp Val His Val Gln Asp Gly Arg
 20 25 30
 Ile Val Ala Leu Gly Val His Ala Glu Ser Val Pro Pro Pro Ala Asp
 35 40 45
 Gln Val Ile Asp Ala Arg Gly Lys Val Val Leu Pro Gly Phe Ile Asn
 50 55 60
 Ala His Thr His Val Asn Gln Ile Leu Leu Arg Gly Gly Pro Ser His
 65 70 75 80
 Gly Arg Gln Phe His Asp Trp Leu Phe Asn Val Val Tyr Pro Gly Gln
 85 90 95
 Lys Ala Met Arg Pro Glu Asp Val Ala Val Ala Val Arg Leu Tyr Cys
 100 105 110
 Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn Ala Asp
 115 120 125

Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val Tyr Gly
 130 135 140
 Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp Arg Met
 145 150 155 160
 Asp Gly Arg Ile Gln Gly Tyr Val Asp Ala Leu Lys Ala Arg Ser Pro
 165 170 175
 Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val Ala Lys Asp
 180 185 190
 Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly Gly Arg
 195 200 205
 Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr Val Glu
 210 215 220
 Gly Met Arg Trp Ala Gln Ala Phe Ala Arg Asp Arg Ala Val Met Trp
 225 230 235 240
 Thr Leu His Met Ala Glu Ser Asp His Asp Gly Arg Ile His Gly Met
 245 250 255
 Ser Pro Ala Glu Tyr Met Glu Cys Tyr Gly Leu Leu Asp Glu Arg Leu
 260 265 270
 Gln Val Ala His Cys Val Tyr Phe Asp Arg Lys Asp Val Arg Leu Leu
 275 280 285
 His Arg His Asn Val Lys Val Ala Ser Gln Val Val Ser Asn Ala Tyr
 290 295 300
 Leu Gly Ser Gly Val Ala Pro Val Pro Glu Met Val Glu Arg Gly Met
 305 310 315 320
 Ala Val Gly Ile Gly Thr Asp Asn Gly Asn Ser Asn Asp Ser Val Asn
 325 330 335
 Met Ile Gly Asp Met Lys Phe Met Ala His Ile His Arg Ala Val His
 340 345 350
 Arg Asp Ala Asp Val Leu Thr Pro Glu Lys Ile Leu Glu Met Ala Thr
 355 360 365
 Ile Asp Gly Ala Arg Ser Leu Gly Met Asp His Glu Ile Gly Ser Ile
 370 375 380
 Glu Thr Gly Lys Arg Ala Asp Leu Ile Leu Leu Asp Leu Arg His Pro
 385 390 395 400
 Gln Thr Thr Pro His His His Leu Ala Ala Thr Ile Val Phe Gln Ala
 405 410 415
 Tyr Gly Asn Glu Val Asp Thr Val Leu Ile Asp Gly Asn Val Val Met
 420 425 430

67

Glu Asn Arg Arg Leu Ser Phe Leu Pro Pro Glu Arg Glu Leu Ala Phe
 435 440 445
 Leu Glu Glu Ala Gln Ser Arg Ala Thr Ala Ile Leu Gln Arg Ala Asn
 450 455 460
 Met Val Ala Asn Pro Ala Trp Arg Ser Leu Glu Met Thr Pro Leu Leu
 465 470 475 480
 His Pro Pro Pro Leu Glu Glu Ile Ala Ala Ile Leu Ala Arg Leu Gly
 485 490 495

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Met Gln Thr Leu Ser Ile Gln His Gly Thr Leu Val Thr Met Asp Gln
 1 5 10 15
 Tyr Arg Arg Val Leu Gly Asp Ser Trp Val His Val Gln Asp Gly Arg
 20 25 30
 Ile Val Ala Leu Gly Val His Ala Glu Ser Val Pro Pro Pro Ala Asp
 35 40 45
 Arg Val Ile Asp Ala Arg Gly Lys Val Val Leu Pro Gly Phe Ile Asn
 50 55 60
 Ala His Thr His Val Asn Gln Ile Leu Leu Arg Gly Gly Pro Ser His
 65 70 75 80
 Gly Arg Gln Phe Tyr Asp Trp Leu Phe Asn Val Val Tyr Pro Gly Gln
 85 90 95
 Lys Ala Met Arg Pro Glu Asp Val Ala Val Ala Val Arg Leu Tyr Cys
 100 105 110
 Ala Glu Ala Val Arg Ser Gly Ile Thr Thr Ile Asn Glu Asn Ala Asp
 115 120 125
 Ser Ala Ile Tyr Pro Gly Asn Ile Glu Ala Ala Met Ala Val Tyr Gly
 130 135 140

Glu Val Gly Val Arg Val Val Tyr Ala Arg Met Phe Phe Asp Arg Met
 145 150 155 160
 Asp Arg Arg Ile Gln Gly Tyr Val Asp Ala Leu Lys Ala Arg Ser Pro
 165 170 175
 Gln Val Glu Leu Cys Ser Ile Met Glu Glu Thr Ala Val Ala Lys Asp
 180 185 190
 Arg Ile Thr Ala Leu Ser Asp Gln Tyr His Gly Thr Ala Gly Gly Arg
 195 200 205
 Ile Ser Val Trp Pro Ala Pro Ala Thr Thr Thr Ala Val Thr Val Glu
 210 215 220
 Gly Met Arg Trp Ala Gln Ala Phe Ala Arg Asp Arg Ala Val Met Trp
 225 230 235 240
 Thr Leu His Met Ala Glu Ser Asp His Asp Glu Arg Ile His Gly Met
 245 250 255
 Ser Pro Ala Glu Tyr Met Glu Cys Tyr Gly Leu Leu Asp Glu Arg Leu
 260 265 270
 Gln Val Ala His Cys Val Tyr Phe Asp Arg Lys Asp Ile Arg Leu Leu
 275 280 285
 His Arg His Asn Val Lys Val Ala Ser Gln Ala Val Ser Asn Ala Tyr
 290 295 300
 Leu Gly Ser Gly Val Ala Pro Val Pro Glu Met Val Glu Arg Gly Met
 305 310 315 320
 Ala Val Gly Ile Gly Thr Asp Asn Gly Asn Ser Asn Asp Ser Val Asn
 325 330 335
 Met Ile Gly Asp Met Lys Phe Met Ala His Ile His Arg Ala Val His
 340 345 350
 Arg Asp Ala Asp Val Leu Thr Pro Glu Lys Ile Leu Glu Met Ala Thr
 355 360 365
 Ile Asp Gly Ala Arg Ser Leu Gly Met Asp His Glu Ile Gly Ser Ile
 370 375 380
 Glu Thr Gly Lys Arg Ala Asp Leu Ile Leu Leu Asp Leu Arg His Pro
 385 390 395 400
 Gln Thr Thr Pro His His His Leu Ala Ala Thr Ile Val Phe Gln Ala
 405 410 415
 Tyr Gly Asn Glu Val Asp Thr Val Leu Ile Asp Gly Asn Val Val Met
 420 425 430
 Glu Asn Arg Arg Leu Ser Phe Leu Pro Pro Glu Arg Glu Leu Ala Phe
 435 440 445

69

Leu Glu Glu Ala Gln Ser Arg Ala Thr Ala Ile Leu Gln Arg Ala Asn
450 455 460

Met Val Ala Asn Pro Ala Trp Arg Ser Leu Glu Met Thr Pro Leu Leu
465 470 475 480

His Pro Pro Pro Leu Glu Glu Ile Ala Ala Ile Leu Ala Gln Leu Gly
485 490 495

What Is Claimed Is:

1. A DNA fragment encoding a homolog of atrazine chlorohydrolase and comprising the sequence of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NOS:7-11 and SEQ ID NOS:17-21.
2. A *s*-triazine-degrading protein having at least one amino acid different from the protein of SEQ ID NO:2, wherein the coding region of the nucleic acid encoding the *s*-triazine degrading protein has at least 95% homology to SEQ ID NO:1 and wherein the *s*-triazine-degrading protein has an altered catalytic activity, as compared with the protein having the sequence of SEQ ID NO:2.
3. The protein of Claim 2 wherein the protein is selected from the group consisting of SEQ ID NOS: 5, 6 and 22-26.
4. The protein of Claim 2 wherein the substrate for the *s*-triazine degrading protein is 2-chloro-4-(ethylamino)-6-(isopropylamino)-1,3,5-triazine.
5. The protein of Claim 2 wherein the substrate for the *s*-triazine degrading protein is 2-chloro-4-(ethylamino)-6-(tertiary butyl-amino)-1,3,5-triazine.
6. The protein of Claim 2 wherein the substrate for the *s*-triazine degrading protein is 2,4,6-triamino-*s*-triazine.
7. A protein selected from the group consisting of proteins comprising the amino acid sequences of SEQ ID NOS: 5, 6 and 22-26.
8. A remediation composition comprising a cell producing the protein of Claim 2.

9. The composition of Claim 8, wherein the composition is suitable for treating soil or water.
10. A remediation composition comprising the protein of Claim 2.
11. The composition of Claim 10 wherein the composition is suitable for treating soil or water.
12. The DNA fragment of Claim 1 in an expression vector.
13. The DNA fragment of Claim 12 in a cell.
14. The DNA fragment of Claim 13 wherein the cell is a bacterium.
15. The DNA fragment of Claim 14 wherein the cell is *E. coli*.
16. A DNA fragment having a portion of its nucleic acid sequence as having at least 95% homology to a DNA fragment consisting of position 236 and ending at position 1655 of SEQ ID NO:1, wherein the DNA fragment is capable of hybridizing under stringent conditions to SEQ ID NO:1 and wherein there is at least one amino acid change in the protein encoded by the DNA fragment as compared with SEQ ID NO:2 and wherein the protein encoded by the DNA fragment is capable of dechlorinating at least one *s*-triazine-containing compound and has an enzymatic activity different from the enzymatic activity of the protein corresponding to SEQ ID NO:2.
17. The fragment of Claim 16, wherein the *s*-triazine-containing compound is 2-chloro-4-(ethylamino)-6-(isopropylamino)-1,3,5-triazine.
18. The fragment of Claim 16, wherein the *s*-triazine-containing compound is 2-chloro-4-(ethylamino)-6-(tertiary butyl-amino)-1,3,5-triazine.

19. The fragment of Claim 16, wherein the *s*-triazine containing compound is (2,4,6-triamino-*s*-triazine).
20. The fragment of Claim 16 wherein the enzymatic activity is an improved ability to degrade atrazine.
21. The fragment of Claim 20 wherein the enzymatic activity is a 10-fold improvement in the ability to degrade atrazine.
22. The fragment of Claim 16, wherein the enzymatic activity is an altered substrate.
23. The protein of Claim 2 which is a homotetramer.
24. The protein of Claim 2 bound to an immobilization support.
25. A method for treating a sample comprising an *s*-triazine-containing compound comprising the step of:
 - adding a composition to a sample comprising an *s*-triazine-containing compound, wherein the composition comprises a protein encoded by a gene having at least a portion of the nucleic acid sequence of the gene having at least 95% homology to the sequence beginning at position 236 and ending at position 1655 of SEQ ID NO:1, wherein the gene is capable of hybridizing under stringent conditions to SEQ ID NO:1, wherein there is at least one amino acid change in the protein encoded by the DNA fragment as compared with SEQ ID NO:2 and wherein the protein has an altered catalytic activity as compared to the protein having the amino acid sequence of SEQ ID NO:2.

26. The method of Claim 25 wherein the composition comprises bacteria expressing the protein.
27. The method of Claim 25 wherein the *s*-triazine -containing compound is 2-chloro-4-(ethylamino)-6-(isopropylamino)-1,3,5-triazine.
28. The method of Claim 25 wherein the *s*-triazine-containing compound is 2-chloro-4-(ethylamino)-6-(tertiary butyl-amino)-1,3,5-triazine.
29. The method of Claim 25 wherein the *s*-triazine containing compound is (2,4,6-triamino-*s*-triazine).
30. The method of Claim 25 wherein the protein encoded by the gene is selected from the group consisting of SEQ ID NOS: 5, 6 and 22-26.
31. A method for obtaining homologs of an atrazine chlorohydrolase comprising the steps of:
- obtaining a nucleic acid sequence encoding atrazine chlorohydrolase;
 - mutagenizing the nucleic acid to obtain a modified nucleic acid sequence that encodes for a protein having an amino acid sequence with at least one amino acid change relative to the amino acid sequence of the atrazine chlorohydrolase;
 - screening the proteins encoded by the modified nucleic acid sequence; and
 - selecting proteins with altered catalytic activity as compared to the catalytic activity of the atrazine chlorohydrolase.
32. The method of Claim 31 wherein the atrazine chlorohydrolase nucleic acid sequence is SEQ ID NO:1.

33. The method of Claim 31 wherein the altered catalytic activity is an improved ability to degrade atrazine.

34. The method of Claim 31 wherein the selected proteins have an altered substrate activity.

1/16

1GCGA 4
1 CTCGGGTAACCTTCTTGAGCGCGGCCACAGCAGCCTTGATCATGAAGGCGA 50
5 GCATGGTGACCTTGACGCCGCTCTTTTCGTTCTCTTTGTTGAACTGCACG 54
51 GCATGGTGACCTTGACGCCGCTCTTTTCGTTCTCTTTGTTGAACTGCACG 100
55 CGAAAGGCTTCCAGGTCGGTGATGTCCGCGTCGTCGTGGTTGGTGACGTG 104
101 CGAAAGGCTTCCAGGTCGGTGATGTCCGCGTCGTCGTGGTTGGTGACGTG 150
105 CGGGATGACCACCCAGTTGCGGTGCAGGTTTTTCGATGGCATAATATCTG 154
151 CGGGATGACCACCCAGTTGCGGTGCAGGTTTTTCGATGGCATAATATCTG 200
155 CGTTGCGACGTGTAACACACTATTGGAGACATATCATGCAAACGCTCAGC 204
201 CGTTGCGACGTGTAACACACTATTGGAGACATATCATGCAAACGCTCAGC 250
205 ATCCAGCACGGTACCCTCGTCACGATGGATCAGTACCGCAGAGTCCTTGG 254
251 ATCCAGCACGGTACCCTCGTCACGATGGATCAGTACCGCAGAGTCCTTGG 300
255 GGATAGCTGGGTTACGTGCAGGATGGACGGATCGTCGCGCTCGGAGTGC 304
301 GGATAGCTGGGTTACGTGCAGGATGGACGGATCGTCGCGCTCGGAGTGC 350
305 ACGCCGAGTCGGTGCCTCCGCCAGCGGATCGGGTGATCGATGCACGCGGC 354
351 ACGCCGAGTCGGTGCCTCCGCCAGCGGATCGGGTGATCGATGCACGCGGC 400
355 AAGGTCGTGTTACCCGGTTTCATCAATGCCCACACCCATGTGAACCAGAT 404
401 AAGGTCGTGTTACCCGGTTTCATCAATGCCCACACCCATGTGAACCAGAT 450
405 CCTCCTGCGCGGAGGGCCCTCGCACGGGCGTCAATTCTATGACTGGCTGT 454
451 CCTCCTGCGCGGAGGGCCCTCGCACGGGCGTCAATTCTATGACTGGCTGT 500
455 TCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGACGTAGCG 504
501 TCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGACGTAGCG 550
505 GTGGCGGTGAGGTTGTATTGTGCGGAAGCTGTGCGCAGCGGGATTACGAC 554
551 GTGGCGGTGAGGTTGTATTGTGCGGAAGCTGTGCGCAGCGGGATTACGAC 600
555 GATCAACGAAAACGCCGATTTCGGCCATCTACCCAGGCAACATCGAGGCCG 604
601 GATCAACGAAAACGCCGATTTCGGCCATCTACCCAGGCAACATCGAGGCCG 650

Fig. 1A

SUBSTITUTE SHEET (RULE 2B)

2/16

605 CGATGGCGGTCTATGGTGAGGTGGGTGTGAGGGTCGTCTACGCCCGCATG 654
|||||
651 CGATGGCGGTCTATGGTGAGGTGGGTGTGAGGGTCGTCTACGCCCGCATG 700
655 TTCTTTGATCGGATGGACGGGCGCATTCAAGGGTATGTGGACGCCTTGAA 704
|||||
701 TTCTTTGATCGGATGGACGGGCGCATTCAAGGGTATGTGGACGCCTTGAA 750
705 GGCTCGCTCTCCCCAAGTCGAACTGTGCTCGATCATGGAGGGAACGGCTG 754
|||||
751 GGCTCGCTCTCCCCAAGTCGAACTGTGCTCGATCATGGAGGAAACGGCTG 800
755 TGGCCAAAGATCGGATCACAGCCCCTGTCAGATCAGTATCATGGCACGGCA 804
|||||
801 TGGCCAAAGATCGGATCACAGCCCCTGTCAGATCAGTATCATGGCACGGCA 850
805 GGAGGTCTGATATCAGTTTGGCCCGCTCCTGCCACTACCACGGCGGTGAC 854
|||||
851 GGAGGTCTGATATCAGTTTGGCCCGCTCCTGCCACTACCACGGCGGTGAC 900
855 AGTTGAAGGAATGCGATGGGCACAAGCCTTCGCCCCGTGATCGGGCGGTAA 904
|||||
901 AGTTGAAGGAATGCGATGGGCACAAGCCTTCGCCCCGTGATCGGGCGGTAA 950
905 TGTGGACGCTTCACATGGCGGAGAGCGATCATGATGAGCGGATTCATGGG 954
|||||
951 TGTGGACGCTTCACATGGCGGAGAGCGATCATGATGAGCGGATTCATGGG 1000
955 ATGAGTCCCGCCGAGTACATGGAGTGTTACGGACTCTTGGATGAGCGTCT 1004
|||||
1001 ATGAGTCCCGCCGAGTACATGGAGTGTTACGGACTCTTGGATGAGCGTCT 1050
1005 GCAGGTCTGCGCATTGCGTGTACTTTGACCGGAAGGATGTTGCGCTGCTGC 1054
|||||
1051 GCAGGTCTGCGCATTGCGTGTACTTTGACCGGAAGGATGTTGCGCTGCTGC 1100
1055 ACCGCCACAATGTGAAGGTCGCGTCGCAGGTTGTGAGCAATGCCTACCTC 1104
|||||
1101 ACCGCCACAATGTGAAGGTCGCGTCGCAGGTTGTGAGCAATGCCTACCTC 1150
1105 GGCTCAGGGGTGGCCCCCGTGCCAGAGATGGTGGAGCGCGGCATGGCCGT 1154
|||||
1151 GGCTCAGGGGTGGCCCCCGTGCCAGAGATGGTGGAGCGCGGCATGGCCGT 1200
1155 GGGCATTGGAACAGATAACGGGAATAGTAATGACTCCGTAAACATGATCG 1204
|||||
1201 GGGCATTGGAACAGATAACGGGAATAGTAATGACTCCGCAAACATGATCG 1250
1205 GAGACATGAAGTTTATGGCCCATATTCACCGCGCGGTGCATCGGGATGCG 1254
|||||
1251 GAGACATGAAGTTTATGGCCCATATTCACCGCGCGGTGCATCGGGATGCG 1300

Fig 1B

SEQUENCE LISTING

3/16

1255 GACGTGCTGACCCCAGAGAAGATTCTTGAAATGGCGACGATCGATGGGGC 1304
|||||
1301 GACGTGCTGACCCCAGAGAAGATTCTTGAAATGGCGACGATCGATGGGGC 1350
1305 GCGTTCGTTGGGAATGGACCACGAGATTGGTTCCATCGAAACCGGCAAGC 1354
|||||
1351 GCGTTCGTTGGGAATGGACCACGAGATTGGTTCCATCGAAACCGGCAAGC 1400
1355 GCGCGGACCTTATCCTGCTTGACCTGCGTCACCCTCAGACGACTCCTCAC 1404
|||||
1401 GCGCGGACCTTATCCTGCTTGACCTGCGTCA.CCTCAGACGACTC..TCA 1447
1405 CATCATTTGGCGGCCACGATCGTGTTTCAGGCTTACGGCAATGAGGTGGA 1454
|||||
1448 CATCATTTGGCGGCCACGATCGTGTTTCAGGCTTACGGCAATGAGGTGGA 1497
1455 CACTGTCCTGATTGACGGAACGTTGTGATGGAGAACCGCCGCTTGAGCT 1504
|||||
1498 CACTGTCCTGATTGACGGAACGTTGTGATGGAGAACCGCCGCTTGAGCT 1547
1505 TTCTTCCCCCTGAACGTGAGTTGGCGTTTCCTTGAGGAAGCGCAGAGCCGC 1554
|||||
1548 TTCTTCCCCCTGAACGTGAGTTGGCGTTTCCTTGAGGAAGCGCAGAGCCGC 1597
1555 GCCACAGCTATTTTGCAGCGGGCGAACATGGTGGCTAACCAGCTTGGGC 1604
|||||
1598 GCCACAGCTATTTTGCAGCGGGCGAACATGGTGGCTAACCAGCTTGGGC 1647
1605 CAGCCTCTAGGAAATGACGCCGTTGCTGCATCCGCCGCCCTTGAGGAAA 1654
|||||
1648 CAGCCTCTAGGAAATGACGCCGTTGCTGCATCCGCCGCCCTTGAGGAAA 1697
1655 TCGCTGCCATCTTGGCGCGGCTCGGATTGGGGGGCGGACATGACCTTGAT 1704
|||||
1698 TCGCTGCCATCTTGGCGCGGCTCGGATTGGGGGGCGGACATGACCTTGAT 1747
1705 GGATACAGAATTGCCATGAATGCGGCACTCCGTCCTTCGCTCGTGTGGA 1754
|||||
1748 GGATACAGAATTGCCATGAATGCGGCACTCCGTCCTTCGCTCGTGTGGA 1797
1755 ATCGTTGGTAGGTGAGGGTCGACTGCGGGCGCCAGCTTCCCGAAGAAGTG 1804
|||||
1798 ATCGTTGGTAGGTGAGGGTCGACTGCGGGCGCCAGCTTCCCGAAGAGGTG 1847
1805 AAAG..... 1808
|||||
1848 AAAGGCCCGAG 1858

Fig. 1C

4/16

1GAGCGCCGCCACAGCAGCCTTGATCATGAAGGCCGA 35
 1 CTCGGGTAACCTTCTTGAGCGCGGCCACAGCAGCCTTGATCATGAAGGCCGA 50
 36 GCATGGTGACCTTGACGCCGCTCTTTTCGTTCTCTTTGTTGAACTGCACG 85
 51 GCATGGTGACCTTGACGCCGCTCTTTTCGTTCTCTTTGTTGAACTGCACG 100
 86 CGAAAGGCTTCCAGGTCGGTGATGTCCGCGTCGTCGTGGTTGGTGACGTG 135
 101 CGAAAGGCTTCCAGGTCGGTGATGTCCGCGTCGTCGTGGTTGGTGACGTG 150
 136 CGGGATGACCACCCAGTTGCGGTGCAGGTTTTTTCGATGGCGTAATATCTG 185
 151 CGGGATGACCACCCAGTTGCGGTGCAGGTTTTTTCGATGGCATAATATCTG 200
 186 CGTTGCGACGTGTAACACACTATTGGAGACATATCATGCAAACGCTCAGC 235
 201 CGTTGCGACGTGTAACACACTATTGGAGACATATCATGCAAACGCTCAGC 250
 236 ATCCAGCACGGTACCCCTCGTCACGATGGATCAGTACCGCAGAGTCCTTGG 285
 251 ATCCAGCACGGTACCCCTCGTCACGATGGATCAGTACCGCAGAGTCCTTGG 300
 286 GGATAGCTGGGTTACAGTGCAGGATGGACGGATCGTCGCGCTCGGAGTGC 335
 301 GGATAGCTGGGTTACAGTGCAGGATGGACGGATCGTCGCGCTCGGAGTGC 350
 336 ACGCCGAGTCGGTGCTCCGCCAGCGGATCGGGTGATCGATGCACGCGGC 385
 351 ACGCCGAGTCGGTGCTCCGCCAGCGGATCGGGTGATCGATGCACGCGGC 400
 386 AAGGTCGTGTTACCCGGTTTTATCAATGCCACACCCATGTGAACCAGAT 435
 401 AAGGTCGTGTTACCCGGTTTTATCAATGCCACACCCATGTGAACCAGAT 450
 436 CCTCCTGCGCGGAGGGCCCTCGCACGGGCGTCAATTCTATGACTGGCTGT 485
 451 CCTCCTGCGCGGAGGGCCCTCGCACGGACGTCAATTCTATGACTGGCTGT 500
 486 TCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGACGTAGCG 535
 501 TCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGACGTAGCG 550
 536 GTGGCGGTGAGGTTGTATTGTGCGGAAGCTGTGCGCAGCGGGATTACGAC 585
 551 GTGGCGGTGAGGTTGTATTGTGCGGAAGCTGTGCGCAGCGGGATTACGAC 600
 586 GATCAACGAAAACGCCGATTTCGGCCATCTACCCAGGCAACATCGAGGCCG 635
 601 GATCAACGAAAACGCCGATTTCGGCCATCTACCCAGGCAACATCGAGGCCG 650

Fig. 2A

5/16

636 CGATGGCGGTCTATGGTGAGGTGGGTGTGAGGGTCGTCTACGCCCCGCATG 685
|||||
651 CGATGGCGGTCTATGGTGAGGTGGGTGTGAGGGTCGTCTACGCCCCGCATG 700
686 TTCTTTGATCGGATGGACGGGCGCATTCAAGGGTATGTGGACGCCTTGAA 735
|||||
701 TTCTTTGATCGGATGGACGGGCGCATTCAAGGGTATGTGGACGCCTTGAA 750
736 GGCTCGCTCTCCCCAAGTCGAACTGTGCTCGATCATGGAGGAAACGGCTG 785
|||||
751 GGCTCGCTCTCCCCAAGTCGAACTGTGCTCGATCATGGAGGAAACGGCTG 800
786 TGGCCAAAGATCGGATCACAGCCCTGTCAGATCAGTATCATGGCACGGCA 835
|||||
801 TGGCCAAAGATCGGATCACAGCCCTGTCAGATCAGTATCATGGCACGGCA 850
836 GGAGGTCGTATATCAGTTTGGCCCGCTCCTGCCACTACCACGGCGGTGAC 885
|||||
851 GGAGGTCGTATATCAGTTTGGCCCGCTCCTGCCACTACCACGGCGGTGAC 900
886 AGTTGAAGGAATGCGATGGGCACAAGCCTTCGCCCCGTGATCGGGCGGTAA 935
|||||
901 AGTTGAAGGAATGCGATGGGCACAAGCCTTCGCCCCGTGATCGGGCGGTAA 950
936 TGTGGACGCTTCACATGGCGGAGAGCGATCATGATGAGCGGATTCATGGG 985
|||||
951 TGTGGACGCTTCACATGGCGGAGAGCGATCATGATGAGCGGATTCATGGG 1000
986 ATGAGTCCCGCCGATTACATGGAGTGTTACGGACTCTTGGATGAGCGTCT 1035
|||||
1001 ATGAGTCCCGCCGAGTACATGGAGTGTTACGGACTCTTGGATGAGCGTCT 1050
1036 GCAGGTCGCGCATTGCGTGTACTTTGACCGGAAGGATGTTGCGCTGCTGC 1085
|||||
1051 GCAGGTCGCGCATTGCGTGTACTTTGACCGGAAGGATGTTGCGCTGCTGC 1100
1086 ACCGCCACAATGTGAAGGTCGCGTCGCAGGTTGTGAGCAATGCCTACCTC 1135
|||||
1101 ACCGCCACAATGTGAAGGTCGCGTCGCAGGTTGTGAGCAATGCCTACCTC 1150
1136 GGCTCAGGGGTGGCCCCCGTGCCAGAGATGGTGGAGCGCGGCATGGCCGT 1185
|||||
1151 GGCTCAGGGGTGGCCCCCGTGCCAGAGATGGTGGAGCGCGGCATGGCCGT 1200
1186 GGGCATTGGAACAGATAACGGGAATAGTAATGACTCCGTAAACATGATCG 1235
|||||
1201 GGGCATTGGAACAGATAACGGGAATAGTAATGACTCCGCAAACATGATCG 1250
1236 GAGACATGAAGTTTATGGCCCATATTACCGCGCGGTGCATCGGGATGCG 1285
|||||
1251 GAGACATGAAGTTTATGGCCCATATTACCGCGCGGTGCATCGGGATGCG 1300

Fig. 2B

SUBSTITUTE SHEET (RULE 2A)

6/16

1286 GACGTGCTGACCCCAGAGAAGATTCTTGAAATGGCGACGATCGATGGGGC 1335
|||||
1301 GACGTGCTGACCCCAGAGAAGATTCTTGAAATGGCGACGATCGATGGGGC 1350
|||||
1336 GCGTTCGTTGGGGATGGACCACGAGATTGGTTCCATCGAAACCGGCAAGC 1385
|||||
1351 GCGTTCGTTGGGAATGGACCACGAGATTGGTTCCATCGAAACCGGCAAGC 1400
|||||
1386 GCGCGGACCTTATCCTGCTTGACCTGCGTCACCCTCAGACGACTCCTCAC 1435
|||||
1401 GCGCGGACCTTATCCTGCTTGACCTGCGTCA.CCTCAGACGACTC..TCA 1447
|||||
1436 CATCATTTGGCGGCCACGATCGTGTTTCAGGCTTACGGCAATGAGGTGGA 1485
|||||
1448 CATCATTTGGCGGCCACGATCGTGTTTCAGGCTTACGGCAATGAGGTGGA 1497
|||||
1486 CACTGTCTTGATTGACGGAAACGTTGTGATGGAGAACCGCCGCTTGAGCT 1535
|||||
1498 CACTGTCTTGATTGACGGAAACGTTGTGATGGAGAACCGCCGCTTGAGCT 1547
|||||
1536 TTCTTCCCCCTGAACGTGAGTTGGCGTTCCTTGAGGAAGCGCAGAGCCGC 1585
|||||
1548 TTCTTCCCCCTGAACGTGAGTTGGCGTTCCTTGAGGAAGCGCAGAGCCGC 1597
|||||
1586 GCCACAGCTATTTTGCAGCGGGCGAACATGGTGGCTAACCAGCTTGGCG 1635
|||||
1598 GCCACAGCTATTTTGCAGCGGGCGAACATGGTGGCTAACCAGCTTGGCG 1647
|||||
1636 CAGCCTCTAGGAAATGACGCCGTTGCTGCATCCGCCGCCCTTGAGGAAA 1685
|||||
1648 CAGCCTCTAGGAAATGACGCCGTTGCTGCATCCGCCGCCCTTGAGGAAA 1697
|||||
1686 TCGTGCCATCTTGGCGCGGCTCGGATTGGGGGGCGGACATGACCTTGAT 1735
|||||
1698 TCGTGCCATCTTGGCGCGGCTCGGATTGGGGGGCGGACATGACCTTGAT 1747
|||||
1736 GGATACAGAATTGCCATGAATGCGGCACTTCCGTCCTTCGCTCGTGTGGA 1785
|||||
1748 GGATACAGAATTGCCATGAATGCGGCACTTCCGTCCTTCGCTCGTGTGGA 1797
|||||
1786 ATCGTTGGTAGGTGAGGGTCGACTGCGGGCGCCAGCTTCCCGAAGAAGTG 1835
|||||
1798 ATCGTTGGTAGGTGAGGGTCGACTGCGGGCGCCAGCTTCCCGAAGAGGTG 1847
|||||
1836 AAAGGCCCGAG 1846
|||||
1848 AAAGGCCCGAG 1858

Fig. 2C

SUBSTITUTE SHEET (RULE 20)

7/16

```

1 .....ASMVTLTPLFSFSLNCTRKASRSVMSASSWLVT 35
  |||
1 SGNFLSAATAALIMKASMVTLTPLFSFSLNCTRKASRSVMSASSWLVT 50
  |||
36 GMTTQLRCRFFDGIISALRRVTHYWRHIMQTL SIQHGT LV TMDQYRRVLG 85
  |||
51 GMTTQLRCRFFDGIISALRRVTHYWRHIMQTL SIQHGT LV TMDQYRRVLG 100
  |||
86 DSWVHVQDGRIVALGVHAESVPPPADRVIDARGKVVLPGFINAHTHVNQI 135
  |||
101 DSWVHVQDGRIVALGVHAESVPPPADRVIDARGKVVLPGFINAHTHVNQI 150
  |||
136 LLRGGPSHGRQFYDWLFNVVYPGQKAMPEDVAVAVRLYCAEAVRSGITT 185
  |||
151 LLRGGPSHGRQFYDWLFNVVYPGQKAMPEDVAVAVRLYCAEAVRSGITT 200
  |||
186 INENADSAIYPGNIEAAMAVYGEVGV RVVYARMFFDRMDGRIQGYVDALK 235
  |||
201 INENADSAIYPGNIEAAMAVYGEVGV RVVYARMFFDRMDGRIQGYVDALK 250
  |||
236 ARSPQVELCSIMEGTAVAKDRITALSDQYHG TAGGRISVWPAPATTTAVT 285
  |||
251 ARSPQVELCSIMEETAVAKDRITALSDQYHG TAGGRISVWPAPATTTAVT 300
  |||
286 VEGMRWAQAFARDRAVMWTLHMAESDHDERIHGMSPAEYMECYGLLDERL 335
  |||
301 VEGMRWAQAFARDRAVMWTLHMAESDHDERIHGMSPAEYMECYGLLDERL 350
  |||
336 QVAHC VYFDRKDVRLLHRHNVKVASQVVS NAYLGSGVAPVPEMVERGMAV 385
  |||
351 QVAHC VYFDRKDVRLLHRHNVKVASQVVS NAYLGSGVAPVPEMVERGMAV 400
  |||
386 GIGTDNGNSNDSVNMIGDMKFMAHIHRAVHRDADVLTPEKILEMATIDGA 435
  |||
401 GIGTDNGNSNDSANMIGDMKFMAHIHRAVHRDADVLTPEKILEMATIDGA 450
  |||
436 RSLGMDHEIGSIETGKRADLILLDLRHPQTTPHHHLAATIVFQAYGNEVD 485
  |||
451 RSLGMDHEIGSIETGKRADLILLDLRHLRRLS.HHLAATIVFQAYGNEVD 499
  |||
486 TVLIDGNVVMENRRLSFLPPERELAFLEEAQSRATAILQRANMVANPAWR 535
  |||
500 TVLIDGNVVMENRRLSFLPPERELAFLEEAQSRATAILQRANMVANPAWR 549
  |||
536 SL*EMTPLLHPPPLEEIAAILARLGLGGGHDLDGYRIAMNAALPSFARVE 585
  |||
550 SL*EMTPLLHPPPLEEIAAILARLGLGGGHDLDGYRIAMNAALPSFARVE 599
  |||
586 SLVGEGRRLRAPASRRSE... 602
  |||
600 SLVGEGRRLRAPASRRGERPE 619
  |||

```

Fig. 3

SUBSTITUTE SHEET (RULE 28)

8/16

```

1 .....SAATAALIMKASMTLTPLFSFSLNCTRKASRSVMSASSWLVT 45
  |||||||||||||||||||||||||||||||||||||||
1 SGNFLSAATAALIMKASMTLTPLFSFSLNCTRKASRSVMSASSWLVT 50
  |||||||||||||||||||||||||||||||||||||||
46 GMTTQLRCRFFDGVISALRRVTHYWRHIMQTL SIQHGT LVTMDQYRRVLG 95
  |||||||||||||:|||||||||||||||||||||||
51 GMTTQLRCRFFDGIISALRRVTHYWRHIMQTL SIQHGT LVTMDQYRRVLG 100
  |||||||||||||:|||||||||||||||||||||||
96 DSWVHVQDGRIVALGVHAESVPPPADRVIDARGKVVLPGFINAHTHVNQI 145
  |||||||||||||||||||||||||||||||||||||||
101 DSWVHVQDGRIVALGVHAESVPPPADRVIDARGKVVLPGFINAHTHVNQI 150
  |||||||||||||||||||||||||||||||||||||||
146 LLRGGPSHGRQFYDWLFNVVYPGQKAMRPEDVAVAVRLYCAEAVRSGITT 195
  |||||||||||||||||||||||||||||||||||||||
151 LLRGGPSHGRQFYDWLFNVVYPGQKAMRPEDVAVAVRLYCAEAVRSGITT 200
  |||||||||||||||||||||||||||||||||||||||
196 INENADSAIYPGNI EAAMAVYGEVGV RVVYARMFFDRMDGRIQGYVDALK 245
  |||||||||||||||||||||||||||||||||||||||
201 INENADSAIYPGNI EAAMAVYGEVGV RVVYARMFFDRMDGRIQGYVDALK 250
  |||||||||||||||||||||||||||||||||||||||
246 ARSPQVELCSIMEETAVAKDRITALSDQYHGTAGGRISVWPAPATTTAVT 295
  |||||||||||||||||||||||||||||||||||||||
251 ARSPQVELCSIMEETAVAKDRITALSDQYHGTAGGRISVWPAPATTTAVT 300
  |||||||||||||||||||||||||||||||||||||||
296 VEGMRWAQAFARDRAVMWTLHMAESDHDERIHGMSPADYMECYGLLDERL 345
  |||||||||||||:|||||||||||||||||||||||
301 VEGMRWAQAFARDRAVMWTLHMAESDHDERIHGMSPA EYMECYGLLDERL 350
  |||||||||||||:|||||||||||||||||||||||
346 QVAHCVFYDRKDVRLLRHNVKVASQVVS NAYLGSGVAPVPEMVERGMV 395
  |||||||||||||||||||||||||||||||||||||||
351 QVAHCVFYDRKDVRLLRHNVKVASQVVS NAYLGSGVAPVPEMVERGMV 400
  |||||||||||||||||||||||||||||||||||||||
396 GIGTDNGNSNDSVNMIGDMKFMAHIHRAVHRDADVLTPEKILEMATIDGA 445
  |||||||||||||.|||||||||||||||||||||||
401 GIGTDNGNSNDSANMIGDMKFMAHIHRAVHRDADVLTPEKILEMATIDGA 450
  |||||||||||||.|||||||||||||||||||||||
446 RSLGMDHEIGSIETGKRADLILLDLRHPQTTPHHHLAATIVFQAYGNEVD 495
  |||||||||||||||||||||||||.|||||||||||
451 RSLGMDHEIGSIETGKRADLILLDLRHLRLS.HHLAATIVFQAYGNEVD 499
  |||||||||||||||||||||.|||||||||||
496 TVLIDGNVVMENRRLSFLPPERELAFLEEAQSRATAILQRANMVANPAWR 545
  |||||||||||||||||||||||||||||||||||||||
500 TVLIDGNVVMENRRLSFLPPERELAFLEEAQSRATAILQRANMVANPAWR 549
  |||||||||||||||||||||||||||||||||||||||
546 SL*EMTPLLHPPPLEEIAAILARLGLGGGHDLDGYRIAMNAALPSFARVE 595
  |||||||||||||||||||||||||||||||||||||||
550 SL*EMTPLLHPPPLEEIAAILARLGLGGGHDLDGYRIAMNAALPSFARVE 599
  |||||||||||||||||||||||||||||||||||||||
596 SLVGEGRLRAPASRRSERPE 615
  |||||||||||||:|||||
600 SLVGEGRLRAPASRRGERPE 619

```

Fig. 4

SUBSTITUTE SHEET (RULE 28)

545 CGGTATCGGGGAATTNTTTGAGCGCGGCCACAGCAGCCNTGATCATGAAGG 496
 1 ...CTCGGGTAACTTCTTGAGCGCGGCCACAGCAGCCTTGATCATGAAGG 47
 495 CGAGCATGGTGACCTNGACGCCGTTNTTTTNGTTNTTTTTTGTGAACTGC 446
 48 CGAGCATGGTGACCTTGACGCCGCTCTTTTCGTTCTCTTTGTGAACTGC 97
 445 ACGCGAAAGG.TTCCAGGTCGGTGATGTCCGCGTCGTCGTGGTTGGTGAC 397
 98 ACGCGAAAGGCTTCCAGGTCGGTGATGTCCGCGTCGTCGTGGTTGGTGAC 147
 396 GTGCGGGATGACCACCCAGNTGCGGTGCAGGTTTTTTCGATGGCATAATAT 347
 148 GTGCGGGATGACCACCCAGTTGCGGTGCAGGTTTTTTCGATGGCATAATAT 197
 346 CTGCGTTGCGACGTGTAACACACTANTGGAGACATATCATGCAAACGCTC 297
 198 CTGCGTTGCGACGTGTAACACACTATTGGAGACATATCATGCAAACGCTC 247
 296 AGCATCCAGCACGGTACCCTCGTCACGATGGATCAGTACCGCAGAGTCCT 247
 248 AGCATCCAGCACGGTACCCTCGTCACGATGGATCAGTACCGCAGAGTCCT 297
 246 TGGGGATAGNTGGGTTACAGTGCAGGATGGACGGATCGTCGCGCTCGGAG 197
 298 TGGGGATAGCTGGGTTACAGTGCAGGATGGACGGATCGTCGCGCTCGGAG 347
 196 TGCACGCCGAGTCGGTGCCTCCGCCAGCGGATCGGGTGATCGATGCACGC 147
 348 TGCACGCCGAGTCGGTGCCTCCGCCAGCGGATCGGGTGATCGATGCACGC 397
 146 GGCAAGGTCGTGTTACCCGGTTTCATCAATGCCACACCCATGTGAACCA 97
 398 GGCAAGGTCGTGTTACCCGGTTTCATCAATGCCACACCCATGTGAACCA 447
 96 GATCCTCCTGCGCGGAGGGCCNTCGCACGGGCGTCAATTNTATGACTGGC 47
 448 GATCCTCCTGCGCGGAGGGCCCTCGCACGGACGTCAATTCTATGACTGGC 497
 46 TGTTCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGA.... 1
 498 TGTTCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGACGTA 547

Fig. 5A

1 ...CCTGCGCGAGGGCCCTCCGCACGGGCGTCAATTCTATGACTGGCTGT 47
 451 CCTCCTGCGCGGAGGGCCCTCGCACGGACGTCAATTCTATGACTGGCTGT 500
 48 TCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGACGTANCG 97
 501 TCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGACGTAGCG 550
 98 GTGGCGGTGAGGTTGTATTGTGCGGAAGCTGTGCGCAGCGGGATTACGAC 147
 551 GTGGCGGTGAGGTTGTATTGTGCGGAAGCTGTGCGCAGCGGGATTACGAC 600
 148 GATCAACGAAAACNCCGATTTCGCCATCTACCCAGGCAACATCGAGGCCG 197
 601 GATCAACGAAAACGCCGATTTCGCCATCTACCCAGGCAACATCGAGGCCG 650
 198 CGATGGCGGTCTATGGTGAGGTGGGTGTGAGGGTCGTCTACGCCCGCATG 247
 651 CGATGGCGGTCTATGGTGAGGTGGGTGTGAGGGTCGTCTACGCCCGCATG 700
 248 TTCTTTGATCGGATGGACGGGCGCATTCAAGGGTATGTGGACGCCTTGAA 297
 701 TTCTTTGATCGGATGGACGGGCGCATTCAAGGGTATGTGGACGCCTTGAA 750
 298 GGCTCGCTCTCCCCAAGTCGAACTGTGCTCGATCATGGANGAAACNGCTG 347
 751 GGCTCGCTCTCCCCAAGTCGAACTGTGCTCGATCATGGAGGAAACGGCTG 800
 348 TGGCCAAAGATCGGATCACANCCCTGTCANATCANTATCATGGCACNGCA 397
 801 TGGCCAAAGATCGGATCACAGCCCTGTGAGATCAGTATCATGGCACGGCA 850
 398 NGAGGTCTATATCANTTTGGCCCGCTCCTGCCACTACCACNGCGGTGAC 447
 851 GGAGGTCTATATCAGTTTGGCCCGCTCCTGCCACTACCACGGCGGTGAC 900
 448 ATTTAAANGAATCCATGGGCCA...ACCTCCCCGTGATCCGGCGGTAA 493
 901 AGTTGAAGGAATGCGATGGGCACAAGCCTTCGCCCGTGATCGGGCGGTAA 950
 494 TGTGAC..... 499
 951 TGTGGACGCTTCACATGGCGGAGAGCGATCATGATGAGCGGATTCATGGG 1000

Fig. 5B

11/16

```

360 .....TNGCAGGTTGTGAGCA..TGCTACTTC 336
      |:|||||
1101 ACCGCCACAATGTGAAGGTCGCGTCGCAGGTTGTGAGCAATGCCTACCTC 1150
      |||:|||||
335 GGTTCAGGNGTGGCCCCCGTGCCAGAGATGGTGGAGCGCGGCATGGCCGT 286
      |||:|||||
1151 GGCTCAGGGGTGGCCCCCGTGCCAGAGATGGTGGAGCGCGGCATGGCCGT 1200
      |||:|||||
285 GGGCATTGGAACAGATAACGGGAATAGTAATGACTCCGTAAACATGATCG 236
      |||:|||||
1201 GGGCATTGGAACAGATAACGGGAATAGTAATGACTCCGCAAACATGATCG 1250
      |||:|||||
235 GAGACATGAAGTTTATGGCCCATATTCACCGCGCGGTGCATCGGGATGCG 186
      |||:|||||
1251 GAGACATGAAGTTTATGGCCCATATTCACCGCGCGGTGCATCGGGATGCG 1300
      |||:|||||
185 GACGTGCTGACCCAGAGAAGATTNTTGAAATGGCGACGATCGATGGGGC 136
      |||:|||||
1301 GACGTGCTGACCCAGAGAAGATTCTTGAAATGGCGACGATCGATGGGGC 1350
      |||:|||||
135 GCGTTTCGTTGGGGATGGACCACGAGATTGGTTCCATCGAAACCGGCAAG 86
      |||:|||||
1351 GCG.TTCGTTGGGAATGGACCACGAGATTGGTTCCATCGAAACCGGCAAG 1399
      |||:|||||
85 CGCGCGGACCTTATCCTGCTTGACCTGCGTCACCCTCAGACGACTCCTCA 36
      |||:|||||
1400 CGCGCGGACCTTATCCTGCTTGACCTGCGTCA.CCTCAGACGACTC..TC 1446
      |||:|||||
35 CCATCATTTGGCGGCCACGATCGTGTTTCAGGCTT..... 1
      |||:|||||
1447 ACATCATTTGGCGGCCACGATCGTGTTTCAGGCTTACGGCAATGAGGTGG 1496

```

Fig. 5C

12/16

1CGGCCACGATCGTGTTCAGGCTTACGGCAATGAGGTGGACAC 43
|||||
1451 CATTTGGCGGCCACGATCGTGTTCAGGCTTACGGCAATGAGGTGGACAC 1500
44 TGCCTGATTGACGGAAACGTTGTGATGGAGAACCGCCGCTTGAGCTTTC 93
|||||
1501 TGCCTGATTGACGGAAACGTTGTGATGGAGAACCGCCGCTTGAGCTTTC 1550
94 TTCCCCCTGAACGTGAGTTGGCGTTCCTTGAGGAAGCGCAGAGCCGCGCC 143
|||||
1551 TTCCCCCTGAACGTGAGTTGGCGTTCCTTGAGGAAGCGCAGAGCCGCGCC 1600
144 ACAGCTATTTTGCATCGGGCGAAACATGGTGGCTAACCCAGCTTGGCGCA 193
|||||
1601 ACAGCTATTTTGCAGCGGGCG.AACATGGTGGCTAACCCAGCTTGGCGCA 1649
194 GCCTCTAGGAAATGACGCCGTTGCTGCATCCGCCGCCCCCTTGAGGAAATC 243
|||||
1650 GCCTCTAGGAAATGACGCCGTTGCTGCATCCGCCGCCCCCTTGAGGAAATC 1699
244 GCTGCCATCTTGGCGCGGCTCGGATTGGGGGGCGGACATGACCTTGATGG 293
|||||
1700 GCTGCCATCTTGGCGCGGCTCGGATTGGGGGGCGGACATGACCTTGATGG 1749
294 ATACAGAATTGCCATGAATGCGGCACTTCCGTCCTTCGCTCGTGTGGAAT 343
|||||
1750 ATACAGAATTGCCATGAATGCGGCACTTCCGTCCTTCGCTCGTGTGGAAT 1799
344 CGTTGGTAGGTGAGGGTCGACTGCGGGCGCCAGCTTCCCGAAGAGGTGAA 393
|||||
1800 CGTTGGTAGGTGAGGGTCGACTGCGGGCGCCAGCTTCCCGAAGAGGTGAA 1849
394 AGCCCGAGGATCCTCTAGAGTCCGATTTTCCGATGTCATCACCGGCGCG 443
|||
1850 AGGCCCGAG..... 1858

Fig. 5D

13/16

1 ...CCTGCGCGGA.GGCCTCCGCACGGGCGTCAATTCTATGACTGGCTGT 46
|||||
451 CCTCCTGCGCGGAGGGCCCTCGCACGGACGTCAATTCTATGACTGGCTGT 500
47 TCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGACGTANCG 96
|||||
501 TCAACGTTGTGTATCCGGGACAAAAGGCGATGAGACCGGAGGACGTAGCG 550
97 GTGGCGGTGAGGTTGTATTGTGCGGAAGCTGTGCGCAGCGGGATTACGAC 146
|||||
551 GTGGCGGTGAGGTTGTATTGTGCGGAAGCTGTGCGCAGCGGGATTACGAC 600
147 GATCAACGAAAACGCCGATTCGGCCATCTACCCAGGCAACATCGAGGCCG 196
|||||
601 GATCAACGAAAACGCCGATTCGGCCATCTACCCAGGCAACATCGAGGCCG 650
197 CGATGGCGGTCTATGGTGAGGTGGGTGTGAGGGTCGTCTACGCCCGCATG 246
|||||
651 CGATGGCGGTCTATGGTGAGGTGGGTGTGAGGGTCGTCTACGCCCGCATG 700
247 TTCTTTGATCGGATGGACGGGCGCATTCAAGGGTATGTGGACGCCTTGAA 296
|||||
701 TTCTTTGATCGGATGGACGGGCGCATTCAAGGGTATGTGGACGCCTTGAA 750
297 GGCTCGCTCTCCCCAAGTCGAACTGTGCTCGATCATGGAGGAAACGGCTG 346
|||||
751 GGCTCGCTCTCCCCAAGTCGAACTGTGCTCGATCATGGAGGAAACGGCTG 800
347 TGGCCAAAGATCGGATCACANCCCTGTCAGATCANTATCATGGCACGGCA 396
|||||
801 TGGCCAAAGATCGGATCACAGCCCTGTCAGATCAGTATCATGGCACGGCA 850
397 NGAGGTCCTATATCANTTTGGCCCGCTCCTGCCACTACCACNGCGGTGAC 446
:|||||
851 GGAGGTCGTATATCAGTTTGGCCCGCTCCTGCCACTACCACGGCGGTGAC 900
447 ATTTNAANGAATTCCATNGGCACAA.CCTTCCCCCGTGATCNGGCGGTAA 495
|:|:|:|
901 AGTTGAAGGAATGCGATGGGCACAAGCCTTCGCCCCTGATCGGGCGGTAA 950
496 TGTNGACCCA..... 505
|:|:|
951 TGTGGACGCTTCACATGGCGGAGAGCGATCATGATGAGCGGATTCATGGG 1000

Fig. 6

14/16




Fig 7a






Fig. 7b

15/16

TERBUYLAZINE DEGRADATION BY SHUFFLED PROTEINS
A7, A11 AND T7 - 1 µg/ml ptn conc
- EFFECT OF METAL IONS



16/16

<u>Bacterium</u>	<u>Translation of PCR amplified DNA sequence</u>					
	79	92				
ADP	SHGRQ	FYDWLFNVVY	PGQKAMRPED	VAVAVRLYCA	EAVRSGITTI	
SG1	PHGRQ	FYDWLFNVLY	PGQKAMRPED	VAVAVRLYCA	EAVRSGITTI	
M91-3	SHGRQ	FYDWLFNVLY	PGQKAMRPED	VAVAVRLYCA	EAVRSGITTI	
J14a	PHGRQ	FYDWLFNVVY	PGQKAMRPED	VAVAVRLYCA	EAVRSGITTI	
38/38	SHGRQ	FYDWLFNVLY	PGQKAMRPED	VAVAVRLYCA	EAVRSGITTI	
Clav.	SHGRQ	FYDWLFNVVY	PGQKAMRPED	VAVAVRLYCA	EAVRSGITTI	
	125			170		
ADP	NE.NADSAIY	PGNIEAAMAV	YGEVGVRVVY	ARMFFDRMDG	RIQGYVDALK	
SG1	NE.NADSAIY	PGNIEAAMAV	YGEVGVRVVY	ARMFFDRMDG	RIQGYVDALK	
M91-3	NE.NADSAIY	PGNIEAAMAV	YGEVGVRVVY	ARMFFDRMDG	RIQGYVDTLK	
J14a	NE.NADSAIY	PGNIEAAMAV	YGEVGVRVVY	ARMFFDRMDG	RIQGYVDALK	
38/38	NENNADSAIY	PGNIEAAMAV	YGEVGVRVVY	ARMFFDRMDG	RIQGYVDTLK	
Clav.	NE.NADSAIY	PGNIEAAMAV	YGEVGVRVVY	ARMFFDRMDG	RIQGYVDALK	
ADP	ARSPQVELCS	IMEETAVAKD	RITALSDQYH	GTAGGRISVW	PAPATTTAVT	
SG1	ARSPQVELCS	IMEETAVAKD	RITALSDQYH	GTAGGRISVW	PAPATTTAVT	
M91-3	ARSPQVELCS	IMEETAVAKD	RITALSDQYH	GTAGGRISVW	PAPATTTAVT	
J14a	ARSPQVELCS	IMEETAVAKD	RITALSDQYH	GTAGGRISVW	PAPATTTAVT	
38/38	ARSPQVELCS	IMEETAVAKD	RITALSDQYH	GTAGGRISVW	PAPATTTAVT	
CLav.	ARSPQVELCS	IMEETAVAKD	RITALSDQYH	GTAGGRISVW	PAPATTTAVT	

Fig. 9

INTERNATIONAL SEARCH REPORT

International Application No

PC 1/US 98/00944

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/55 C12N9/14 C07K16/40 C12Q1/34 C12N11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 97 15675 A (UNIV MINNESOTA) 1 May 1997 see the whole document	1-34
X	--- SOUZA DE M L ET AL: "ATRAZINE CHLOROXYDROLASE FROM PSEUDOMONAS SP. STRAIN ADP: GENE SEQUENCE, ENZYME PURIFICATION, AND PROTEIN CHARACTERIZATION" JOURNAL OF BACTERIOLOGY, vol. 178, no. 16, August 1996, pages 4894-4900, XP000614970 see the whole document --- -/--	1-34

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 May 1998

Date of mailing of the international search report

25/05/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hillenbrand, G

INTERNATIONAL SEARCH REPORT

International Application No

PCI/US 98/00944

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
X	<p>SOUZA DE M L ET AL: "CLONING, CHARACTERIZATION, AND EXPRESSION OF A GENE REGION FROM PSEUDOMONAS SP. STRAIN ADP INVOLVED IN THE DECHLORINATION OF ATRAZINE"</p> <p>APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 61, no. 9, September 1995, pages 3373-3378, XP000616901</p> <p>see the whole document</p>	1-34
A	<p>MANDELBAUM R T ET AL: "ISOLATION AND CHARACTERIZATION OF A PSEUDOMONAS SP. THAT MINERALIZES THE S-TRIAZINE HERBICIDE ATRAZINE"</p> <p>APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 61, no. 4, April 1995, pages 1451-1457, XP000617154</p> <p>see the whole document</p>	1,2,16,25
A	<p>MULBRY W W: "PURIFICATION AND CHARACTERIZATION OF AN INDUCIBLE S-TRIAZINE HYDROLASE FROM RHODOCOCCUS CORALLINUS NRRL B-15444R"</p> <p>APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 60, no. 2, February 1994, pages 613-618, XP000645496</p> <p>see the whole document</p>	1,2,16,25
A	<p>EP 0 141 784 A (CIBA GEIGY AG) 15 May 1985</p> <p>see the whole document</p>	1,2,16,25

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/00944

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9715675	A	01-05-1997	AU 7466996 A	15-05-1997
EP 0141784	A	15-05-1985	AU 3493084 A	09-05-1985
			BR 8405592 A	10-09-1985
			DE 3466668 A	12-11-1987
			JP 60114191 A	20-06-1985
			US 4745064 A	17-05-1988